

THE FLORISTIC RICHNESS OF BUDA AND RÂIOSU MOUNTAINS, FĂGĂRAȘ MASSIF

STANCU Daniela Ileana

Abstract. The present study deals with the floristic richness of Râiosu and Buda Mountains, Făgăraș Massif. Many endemics and relicts of these vegetal groups confirm the oldness of the vegetation of these calcareous mountains; they survived to the extreme catatherme of the glaciations, because of the thermal characteristics of the calcareous substrate. During the glacial periods, Râiosu and Buda Peaks, that are higher than 2,300 m–2,400 m, functioned as an important "nunatak", which enabled the preservation of this rich flora (endemics and relicts).

Keywords: glacial relief, climate oscillations, relicts, calciphile, Făgăraș.

Rezumat. Bogăția floristică a munților Râiosu și Buda, masivul Făgăraș. Lucrarea prezintă bogăția floristică a munților Râiosu și Buda, din masivul Făgăraș. Mulțimea endemitelor și a relictelor vegetale confirmă vechimea vegetației acestor munți calcaroși; ele au supraviețuit catatermelor glaciațiunilor datorită caracteristicilor termice ale substratului calcaros. Faptul că Vârful Râiosu și Buda sunt situate la peste 2.300-2.400 m altitudine a făcut ca acești munți să funcționeze în perioadele glaciare ca un important „nunatak” care a ajutat la păstrarea acestei flore bogate în endemite și relice.

Cuvinte cheie: relief glaciari, oscilații climatice, relice, calcifile, Făgăraș.

INTRODUCTION

The Făgăraș Mountains preserves the most representative glacial landforms of the Southern Carpathians. The unique features of the landscape of the Romanian Carpathians are mainly induced by the extension of the glacial and periglacial relief and by the highest values of the relief intensity within the entire country. This is why, Emm. de Martonne (1907) called the Southern Carpathians the "Transylvanian Alps". As a consequence of their past the Făgăraș Mountains have a lot of glacial traces: cirques, lakes (mountain lakes) and glacial valleys. The northern slopes characterized by a strong declivity, limited the development of the glacier length, while in the South the gentle and large valleys permitted the formation of some (valley) glaciers that are over 7 km long having a sinuous aspect and ramifications, which derive from the suspended tributary valleys (PIȘOTA, 1968).

The glacial cirques, locally called "bucket" have different aspects and complexity. The valleys Capra and Buda are the most extended; they are formed by "complexes of glacial valleys", but the most spectacular valleys can be found on the western side of Buda – Râiosu – Mușeteica Peak. Some of the valleys have small lakes and glacial cirques that were occupied in the past by the tongues of the glaciers.

The altitude of 2,000-2,200 meters where they are situated indicate the fact that they were formed in a most recent glacial era that corresponds to Würm period (VAN CAMPO, 1969).

The glacial cut blade like indents appeared at the intersection of the glacial valleys and cirques, being accentuated by the past periglacial phenomena or by the actual crionival ones. They are common in the relief of the main crest of the southern slope Buda – Râiosu - Mușeteica. The studied mountains are situated in the northern side of Argeș County; they are a part of the southern slope of the Făgăraș Mountains.

Buda Mountain (2,431 m) is situated in the North of Buda Lake, a glacial lake situated at 2,055 m altitude. The lake has a surface of about 0.86 ha, it is triangular, and its maximum depth is 2.2 m (PIȘOTA, 1971). The Buda Valley has its origins in Buda cirque; it flows southwards and crosses a limestone area forming Gegiu Gorges, with hardly accessible walls. The valley of the Buda Stream is 18 km long and it flows in to Vidrar Lake. It has many tributaries including the Râiosu Stream.

The other studied mountain is Râiosu (2,395 m), which is a continuation of the Buda Mountain together with which it forms an almost continuous peak, as long as they are separated only by a little "saddle" called "Polița lui Vodă" or "Vodă's road". This mountain has the shape of a huge pyramid and it presents all the features of a calcareous mountain with abrupt walls and detritus; the dolomite calcareous intercalation is well developed in sericite schists.

Râiosu Lake has a 0.15 ha surface and it depends on the rainy periods (PIȘOTA, 1971). It is situated at the southern side of the massif, East of Buda – Râiosu – Mușeteica summit, at 2,180 m altitude, in the glacial cirque dominated by Mușeteica Peak at West and by Râiosu Peak at East. This is probably the most spectacular landscape from the Făgăraș Mountains. The lake is strongly clogged because the rocks are desegregated and then transported by the masses of snow and ice. In the summer, it is frequently covered by iced snow. As there is no marked path, the access to the lake is very difficult (STANCU, 2005).

MATERIALS AND METHODS

If in the Pleistocene the characters of the climate justified the term periglacial on the territory of Romania, nowadays the region evolves under a crionival modelling regime at altitudes above the timberline. The crionival

processes and the forms resulted presently are connected with the much more intense and extended manifestation from the Pleistocene; some of them continue their evolution, while others stopped, but they may be reactivated mainly due to the characteristics of the climate, as well as to the anthropogenic action.

The alpine level of the Făgăraș Mountains is characterized by the though climate conditions: the average temperatures of the cold months are -8°C and even -11°C , while the average number of winter days is 120-160. There are 220-250 frozen days yearly. The biggest thermal differences are registered in the transition months: April-May and October-November. Record values of the atmospheric rainfalls are registered in the western part of the massif (over 1,300 mm) while on the eastern side the value is 1,200 mm. Over a half of them are solid rainfalls. The average number of days with snow layer is over 200. The strong wind blowing from West could frequently reach 120 km/h.

The longitudinal orientation of the main alpine summit leads to great thermal differences between the shady northern slopes and the sunny ones, where the nebulosity is higher. The Făgăraș Mountains are considered to be the greatest clouds maker of our country. Beside the clouds coming together with the winds from the Atlantic, from the polar seas or from the Mediterranean Basin, the Făgăraș Mountains forms their own clouds. Some of them appear as a consequence of a stronger warming of the air from the southern slope by comparison with the northern one.

The crionival relief is the direct result of the diurnal thermal oscillations, with maximum efficiency in the aforementioned seasons of transition, but mainly in the spring (FLOREA, 1998). The fact that these mountains are entirely crystalline in their structure (but displaying a great petrographic diversity with intercalations of crystalline limestones and marbles) with cracks that permit water circulation, justifies the variety and the frequency of these forms.

RESULTS AND DISCUSSION

Buda (2,431 m) - Râiosu (2,395 m) - Mușeteica (2,448 m) is the only southern summit that displays the aspect of a ridge surrounded by pyramidal peaks. The presence of the crystalline limestone and dolomites, which led to the appearance of various forms (such as needles, sharp crests, towers, short abrupt sectors) explains the crenellated character of this summit with numerous elements of ruiniform calcareous relief. The limestones from the Râiosu and Buda Mountains shelter a series of vegetal calciphile associations which, despite the localization of these mountains, are representative for the Romanian Carpathians.

Many endemics and relicts of these vegetal groups confirm the oldness of the vegetation of this calcareous massif; they survived to the extreme catatherme of the glaciations, because of the thermal characteristics of the calcareous substrate. The absence of a glacial phenomenology from the calcareous massif shows that Râiosu and Buda, which exceed 2,300 m – 2,400 m, in the glacial periods, functioned as an important "nunatak" contributing to the preservation of this rich flora (endemics and relicts) (BĂNĂRESCU & BOȘCAIU, 1973). The oldness of this flora is also confirmed by the reduced capacity of phytocoenotic competition, a fact which conditioned the surviving in groups situated on rocks or on mobile and semi fixed detritus (petrophille).

The flora of these two mountains includes a great variety of phyto-geographical elements determined by the altitude variations (600 – 2,431 m). The Eurasian elements are dominant within this researched territory, followed by the European, Circumpolar, and Central European ones; the Carpathian endemics and the Carpathian-Balkan elements and the Alps-Carpathian elements represent 18% of this mountains flora (BELDIE, 1967).

On the other hand, a clear relationship between the altitude and the frequency of the calciphile species exists. But many of these species could be also found frequently on some basic rocks, especially gabbros. The Mesozoic limestone enclave of the Râiosu Mountain leads to some general conclusions regarding the floral-genetic role of the calciphile elements (BOȘCAIU, 1975).

In MERXMÜLER (1952) opinion, disjunctions between the calciphile species that can be found these days in the northern and southern parts of the Alps, appeared towards the end of the Tertiary age, when there still where large areas of limestone which, these kinds of plants used to grow on an unbroken area. As a result, the calcareous layer that was in contact with the vegetation could play a bigger part than it does today, in the migration of the calciphile plants. The erosion of the different limestone areas has been directed to isolate through crumble of the calciphile plants' area.

According to Malyshev's explanations (1965), the calciphily developed in the context of the drying of the climate because it ensures the compensation of a deficit of humidity during the periods of drought. As a result, the calciphily played an important role during the times of widen climatic changes that coincided with the formation of the complexes high level of the mountain flora. As a consequence of widen climatic tolerance, the calciphily encourages the migration of diverse vegetal populations across big distances, which supports the floral changes between distant mountains. So, it was favoured the extension of the cryophilic species in regions where the climate was dry. At the same time, the optional calciphily of some cryophilic species, became obligatory in the regions with a humid climate, where they can be found only in a calcareous layer.

According to Malyshev, many species the area of which has far-away disjunction are calciphile species. In its turn, the frequency of the calciphile areas seems to be correlated to the allophone elements of the flora. In Malyshev's opinion, in the native floral complexes, the calciphile species are less numerous than the allochthonous ones. Thus, calciphily favoured the floristic exchanges between the floras of several massifs of the alpine system and the arctic flora. The calciphile plants could migrate on long distances under the conditions of a climate that grows arid or cold in catathermes periods.

Taking these into account, it is accountable that the calciphile plants of the Făgăraș Mountains have polygenetic origins, coming from different phytogeographic areas and their migration happened in different phyto historic ages. Despite the huge phytogeographic interest, the chorological origin of these plants stands relative, as we do not always know the whole data concerning a certain area of all these species.

The Carpathian-Balkan species are very frequent both on the meadows from the alpine space and on the calcareous rocks from the forests (NEGREAN & OLTEAN, 1989). They highlight the floral-genetic connections between the flora of this East-Carpathian space and that of the Balkan Mountains, which existed in the Tertiary. Representative species: *Carduus kernerii* ssp. *kernerii*, *Campanula patula* ssp. *abietina*, *C. transsilvanica*, *Alyssum repens*, *Erysimum witmannii*, *Rhododendron myrtifolium*, *Poa media*, *Sesleria rigida*, *Ranunculus crenatus*, *Potentilla aurea* ssp. *chrysocraspeda*, *Saxifraga luteo-viridis*, *Veronica baumgartenii*, *Viola declinata*, *V. dacica* etc.

The taxa *Aconitum toxicum* highlights the link with the Balkan-Illyrian flora.

The existence of floral historical interferences with Anatolia and Caucasus in the past is attested by the presence of the following taxa: *Bruckenthalia spiculifolia*, *Plantago gentianoides*, *Telekia speciosa*, *Carex nigra* ssp. *dacica*.

A series of western and south-western and even Oriental-continental elements were found in this area of research: *Taraxacum nigricans*, *Scrophularia scopolii*, *Isatis tinctoria* ssp. *praecox*.

The results of the floristic analysis confirms the fact that this territory belongs to the Holarctic region, Eurosiberian sub region, Central-European domain, Eastern-Charpatian (Dacian) province, the Southern Carpathian circumscription, the Făgăraș Mountains district (BORZA & BOȘCAIU, 1965).

The impact of the anthropogenic - zoogenous pressure leads to the endangering and even to the extinction of some species or plants communities. Thus, although some species are still not in danger on a large scale, e.g. the national territory, they could be jeopardized in individual counties. These extinctions already represent an alarm because the extinction of some species in a county is followed by its extinction on the entire national territory (BOȘCAIU et al., 1994).

CONCLUSIONS

According to the phytogeographic information we have nowadays, we propose the following chorological classification of the calciphile plants from the Râiosu and Buda Mountains (CIOCÂRLAN, 2000; DIHORU & PÂRVU, 1987; OPREA, 2005).

Boreo-Circumpolar: *Adoxa moschatellina* L., *Artemisia vulgaris* L., *Asplenium scolopendrium* L., *A. ramosum* L., *Campanula rotundifolia* L., *Carex pallescens* L., *C. sylvatica* HUDSON, *Cerastium arvense* L. ssp. *arvense*, *Circaea alpina* L., *Chrysosplenium alternifolium* L., *Clinopodium vulgare* L., *Coeloglossum viride* (L.) HARTMAN, *Eriophorum latifolium* HOPPE, *Equisetum hiemale* L., *E. pratense* EHRH., *Galium aparine* L., *Geum aleppicum* JACQ., *G. rivale* L., *Gnaphalium sylvaticum* L., *Gymnocarpium robertianum* (HOFFM.) NEWMAN, *Hepatica nobilis* SCHREBER, *Juniperus communis* L., *Koeleria macrantha* (LEDEB) SCHULTES, *Milium effusum* L., *Moneses uniflora* (L.) A. GRAY., *Parnassia palustris* L., *Polypodium vulgare* L., *Polystichum braunii* (SPENNER) FÉE, *P. lonchitis* (L.) ROTH, *Pyrola minor* L., *Rubus idaeus* L., *Sagina procumbens* L., *Selaginella selaginoides* (L.) LINK, *Spergularia rubra* (L.) J. et C. PRESL, *Stellaria uliginosa* MURRAY, *Viola biflora* L.

Arctic-Alpine-Circumpolar: *Aster alpinus* L., *Athyrium distentifolium* TAUSCH ex OPIZ, *Bartsia alpina* L., *Carex atrata* L., *Cerastium cerastoides* (L.) BRITTON, *Cystopteris montana* (LAM.) DESV., *Erigeron uniflorus* L., *Kobresia myosuroides* (VILL.) FIORI, *Lloydia serotina* (L.) REICHENB., *Loiseleuria procumbens* (L.) DESV., *Myosotis alpestris* F. W. SCHMID, *Pedicularis verticillata* L., *Potentilla crantzii* (CRANTZ) BECK ex FRITSCH, *Polygonum viviparum* L., *Rhodiola rosea* L., *Salix alpina* SCOP., *Saxifraga hieracifolia* WALDST. & KIT., *Saussurea alpina* (L.) DC., *Silene acaulis* (L.) JACQ.

Alpine-Circumpolar: *Anemone narcissiflora* L., *Carex pyrenaica* WAHLBG., *Dryas octopetala* L., *Hedysarum hedysaroides* (L.) SCHINZ et THELL., *Myosotis alpestris* F. W. SCHMIDT, *Polygonum viviparum* L., *Salix reticulata* L., *Saxifraga oppositifolia* L.

Alpine-Arctic: *Arabis alpina* L., *Ligusticum mutellinoides* (CRANTZ) VILL., *Nigritella nigra* (L.) REICHENB., *Pedicularis oederi* VAHL., *Saxifraga aizoides* L., *S. androsacea* L., *S. paniculata* MILL., *Pinguicula alpina* L.

Alpine-Altaic: *Astragalus australis* (L.) LAM., *Gentiana verna* L. ssp. *verna*, *Leontopodium alpinum* CASS., *Poa supina* SCHRAD., *Saxifraga moschata* WULFEN.

Alpine-Central-European: *Astragalus alpinus* L., *A. australis* (L.) LAM., *Bellardiochloa violacea* (BELLARDI) CHIOV., *Biscutella laevigata* L., *Bupleurum falcatum* L. ssp. *cernuum* (TEN.) ARCANGELI, *Carex sempervirens* VILL., *Cerastium alpinum* L., *Cerintho glabra* MILLER, *Epilobium alpestre* (JACQ.) KROCKER, *Euphrasia salisburgensis* FUNCK., *Helianthemum alpestre* (JACQ.) D.C., *H. nummularium* (L.) MILLER ssp. *grandiflorum* (SCOP.) SCHINZ. & THELL., *Hieracium aurantiacum* L., *H. villosum* JACQ., *Kernera saxatilis* (L.) REICHENB., *Linaria alpina* (L.) MILLER, *Primula halleri* J.F. GMELIN, *Pritzelago alpina* (L.) O. KUNTZE ssp. *brevicaulis* (SPRENGEL) GREUTER et BURDET, *Ranunculus alpestris* L., *R. oreophilus* BIEB., *Rumex scutatus* L., *Salix retusa* L. ssp. *kitaibeliana* WILLD., *Saxifraga retusa* GOUAN, *S. stellaris* L., *Sedum alpestre* VILL., *S. atratum* L., *Silene pusilla* WALDST. & KIT., *Solidago*

virgaurea L. ssp. *minuta* (L.) ARCANGELI, *Thesium alpinum* L., *Trifolium badium* SCHREB., *Valeriana montana* L., *Veronica aphylla* L.

Alpine-Carpathian: *Acinos alpinus* (L.) MOENCH ssp. *alpinus*, *Aconitum tauricum* WULFEN, *Androsace lactea* L., *A. villosa* L. ssp. *arachnoidea* (SCOTT., NYM. et KOTSCH.) NYM., *Armeria alpina* WILLD., *Campanula alpina* JACQ., *Centaurea nervosa* WILLD., *Dianthus glacialis* HAENKE ssp. *glacialis*, *Draba kotschyi* STUR, *Festuca versicolor* TAUSCH, *Galium anysophyllum* VILL., *Gentiana brachyphylla* VILL., *G. utriculosa* L., *Helianthemum alpestre* (JACQ.) D.C., *Nigritella rubra* (WETTST.) K. RICHTER, *Oxytropis halleri* BUNGE ex KOCH, *Polygala alpestris* REICHENB., *Pulsatilla alba* REICHENB., *Scabiosa lucida* VILL., *Senecio carniolicus* WILLD., *Viola alpina* JACQ.

Alpine-Carpathian-Balkan: *Achillea stricta* (KOCH) SCH., *Aconitum degenii* GAYER, *Alyssum repens* BAUMG. ssp. *repens*, *Alnus viridis* (CHAIX) D.C., *Arabis allionii* D.C., *Crepis conyzifolia* (GOUAN) A. KERN., *Dianthus barbatus* L. ssp. *compactus* KIT., *Doronicum columnae* TEN., *Gentianella lutescens* (VALEN.) J. HOLUB, *Laserpitium krapfii* CRANTZ., *Ligusticum mutellina* (L.) CR., *Phyteuma confusum* A. KERNER, *Ranunculus crenatus* WALDST. & KIT., *Senecio subalpinus* KOCH, *Tanacetum corymbosum* (L.) SCHULTZ BIP. ssp. *subcorimbozum* (SCHUR) PAWL., *Trifolium pallescens* SCHREB.

Balkan-Carpathian (incl. Dacian): *Aconitum toxicum* RCHB., *Alyssum repens* BAUMG. ssp. *transsilvanicum* (SCHUR) NYM., *Asperula capitata* KIT. ex SCHULTES, *Bellardiochloa variegata* (LAM.) KERGUELEN, *Bromus barcensis* SIMONKAI, *Bupleurum diversifolium* ROCH., *Centaurea atropurpurea* WALDST. et KIT., *C. kotschyana* HEUFFEL, *Cirsium candelabrum* GRISEB., *Dianthus petraeus* WALDST. et KIT., *Draba lasiocarpa* ROCHEL, *Galium kitaibelianum* SCHULTES et SCHULTES, *Geranium sylvaticum* L. ssp. *caeruleatum* (SCHUR) D. A. WEB., *Gypsophila petraea* (BAUMG.) REICHENB., *Helleborus purpurascens* WALDST. et KIT., *Lathyrus hallersteinii* BAUMG., *Linum extraaxilare* KIT., *Poa cenisia* ALL. ssp. *contracta* NYAR., *Plantago gentianoides* SIBTH. et SM., *Pulmonaria rubra* SCHOTT, *Ranunculus pseudomontanus* SCHUR, *Saxifraga carpatica* RCHB., *S. corymbosa* BLOSS., *S. pedemontana* ALL. ssp. *cymosa* ENGLER, *S. rotundifolia* L. ssp. *heucherifolia* (GRIS. et SCHENCK) CIOCÂRLAN, *Sempervivum marmoreum* GRISEB., *Senecio pappopus* (RCHB.) LESS., *Sesleria bielzii* SCHUR, *S. rigida* HEUFF. ssp. *haynaldiana* SCHUR, *Silene lichenfeldiana* BAUMG., *Symphandra wanneri* (ROCHEL) HEUFFEL, *Thymus balcanus* BORB., *Veronica baumgartenii* ROEMER et SCHULTES.

Balkan-Dacian-Pannonic: *Chamaecytisus hirsutus* (L.) LINK. ssp. *leucotrichus* (SCHUR) LÖVE et LÖVE.

Carpathian (incl. End): *Achillea schurii* SCHUR, *Aconitum moldavicum* HACQ., *Aquilegia transsilvanica* SCHUR, *Athamantha turbith* (L.) BROT. ssp. *hungarica* (BORBAS) TUTIN, *Campanula carpatica* JACQ., *C. transsilvanica* SCHUR, *Cardaminopsis neglecta* (SCHULTES) HAYEK, *Centaurea pinnatifida* SCHUR, *Cerastium arvense* L. ssp. *lerchenfeldianum* (SCHUR) ASCHERSON et GRAEBNER, *C. transsilvanicum* SCHUR, *Chrysosplenium alpinum* SCHUR, *Dentaria glandulosa* W. et K., *Dianthus glacialis* HAENKE ssp. *gelidus* (SCHOTT, NYM. et KOTSCHY) NYM., *D. henteri* HEUFFEL ex GRISEB., *D. petraeus* WALDST. et KIT. ssp. *petraeus*, *D. spiculifolius* SCHUR, *D. tenuifolius* SCHUR, *Erysimum witmanni* ZAWADSKI ssp. *transsilvanicum* (SCHUR) P. W. BALL, *E. wittmannii* ZAM. ssp. *wittmannii*, *Eritrichium nanum* (L.) SCHRADER ssp. *jankae* (SIMK.) JAV., *Festuca rupicola* HEUFFEL ssp. *saxatilis* (SCHUR) RAUSCHERT, *Genista tinctoria* L. ssp. *oligosperma* (ANDRAE) PROD., *Helictotrichon decorum* (JANKA.) HENRARD, *Hepatica transsilvanica* FUSS, *Hesperis moniliformis* SCHUR, *Leucanthemum waldsteini* (SCHULTZ BIP.) POUZAR, *Onobrychis montana* D. C., ssp. *transsilvanica* (SIMONKAI) JAV., *Origanum vulgare* L. var. *barcense* SIMK., *Oxytropis carpatica* UECHTR., *Papaver alpinum* L. ssp. *corona-sancti-stephani* (ZAPAL.) BORZA, *Phyteuma tetramerum* SCHUR, *Poa granitica* BR.-BL. ssp. *disparilis* (NYAR.) NYAR., *Ranunculus carpaticus* HERBICH, *Symphytum cordatum* WALDST. et KIT., *Thymus comosus* HEUFFEL, *T. pulcherimus* SCHUR.

ACKNOWLEDGMENTS

I gratefully acknowledge the assistance and the support of acad. Nicolae Boșcaiu for his helpful comments and for his advices along my work.

REFERENCES

- BĂNĂRESCU P. & BOȘCAIU N. 1973. *Biogeografie. Perspectivă genetică și istorică*. Edit. Științifică. București. 302 pp.
- BELDIE AL. 1967. *Endemismele și elementele dacice din flora Carpaților românești*. Comunicări de botanică la a V-a Consfătuire de geobotanică. București: 113-120.
- BORZA AL. & BOȘCAIU N. 1965. *Introducere în studiul covorului vegetal*. Edit. Academiei Române. București. 340 pp.
- BOȘCAIU N. 1975. *Problemele conservării vegetației alpine și subalpine*. Ocrotirea Naturii București. **19**(1): 17-21.
- BOȘCAIU N., COLDEA GH., HOREANU CL. 1994. *Lista roșie a plantelor vasculare dispărute, periclitare, vulnerabile și rare din România*. Ocrotirea naturii și a mediului înconjurător. **38**(1): 45-56.
- CIOCÂRLAN V. 2000. *Flora ilustrată a României. Pteridophyta et Spermatophyta*. Ediția a doua revăzută și adăugită. Edit. Ceres. București. 1139 pp.
- DIHORU G. & PÂRVU C. 1987. *Plante endemice în flora României*. Edit. Ceres. București. 180 pp.
- FLOREA M. 1998. *Munții Făgărașului. Studiu geomorfologic*. Edit. Foton. Brașov. 114 pp.

- MALYSHEV L. I. 1965. *Flora alpina Montium Sajanensium Orientalium Nauka*. Moscova-Leningrad. 367 pp (In Russian).
- MERXMÜLLER H. 1952. *Untersuchungen zur Sippengliederung und Arealbildung in den Alpen*. München. 105 pp.
- NEGREAN G. & OLTEAN M. 1989. *Endemite și zone endemo-conservatoare din Carpații S-E*. Ocrotirea naturii și a mediului înconjurător. **33**(1): 15-25.
- OPREA A. 2005. *Lista critică a plantelor vasculare din România*. Edit. Universității. "Al. I. Cuza". Iași. 668 pp.
- PIȘOTA I. 1968. *Lacurile glaciare din Munții Rodnei*. Analele Universității București. (Șt. nat.) Geologie-Geografie. **12**(2): 113-124.
- PIȘOTA I. 1971. *Lacurile glaciare din Carpații Meridionali*. Studiu hidrologic. Edit. Academiei. București. 162 pp.
- STANCU DANIELA ILEANA. 2005. *Flora și vegetația Munților Râiosu și Buda, masivul Făgăraș*. Edit. Universității din Pitești. 226 pp.
- VAN CAMPO M. 1969. *Végétation würmienne en France. Données bibliographiques. Hypothèse*. Etudes française sur le quaternaire présentées à l'occasion du VIII-e Congrès International de l'INQUA. Paris: 1-8.

Stancu Daniela Ileana

Argeș County Museum, A. Călinescu Street 44,

Pitești, România

E-mail: stancuileana@yahoo.com

Received: April 29, 2010

Accepted: August 19, 2010