

## CHANGES OF SOME PHYSIOLOGICAL PARAMETERS IN PRUSSIAN CARP (*Carassius auratus gibelio* BLOCH 1782) UNDER THE ACTION OF THE FOLPAN 80 WDG

ZGURSCHI Gabriela, PONEPAL Cristina, MARINESCU Al. Gabriel

**Abstract.** The massive use of xenobiotics in agriculture has stimulated the research of those compounds in the environment. Folpan 80 WDG is a fungicide used on a broad range of phytopathogenic fungi. This study was carried out to analyze the effects of sub lethal and lethal concentrations – from  $25 \times 10^{-5}$ g Folpan 80 WDG to  $1 \times 10^{-3}$ g Folpan 80 WDG /l water on some physiological parameters (oxygen consumption, breathing frequency, number of erythrocytes) on Prussian carp (*Carassius auratus gibelio* BLOCH 1782). The acute and subacute toxicity of Folpan 80 WDG fungicide was evaluated in glass aquaria. The Folpan 80 WDG product, under the concentrations from  $25 \times 10^{-5}$ g Folpan /l water to  $1 \times 10^{-3}$  Folpan 80 WDG g/l water, produced, after one week of immersion, a significant decrease in the fish oxygen consumption. The fungicide changed the fish respiratory rhythm in all investigated concentrations after 14 days of exposure. The number of erythrocytes significantly decreased after 14 days of immersion at fungicide concentrations.

**Keywords:** Prussian carp, Folpan 80 WDG, oxygen consumption, respiratory rhythm, number of erythrocytes.

**Rezumat. Modificările unor parametri fiziologici la caras (*Carassius auratus gibelio* BLOCH 1782) sub acțiunea Folpanului 80 WDG.** Utilizarea masivă a xenobioticelor în agricultură a stimulat cercetarea acestor compuși în mediul înconjurător. Folpan 80 WDG este un fungicid folosit asupra unui spectru larg de ciuperci fitopatogene. Acest studiu a fost efectuat pentru a analiza efectele concentrațiilor subletale și letale - de la  $25 \times 10^{-5}$ g Folpan 80 WDG la  $1 \times 10^{-3}$ g Folpan 80 WDG /l de apă asupra unor parametri fiziologici (consumul de oxigen, frecvența respiratorie, numărul de eritrocite) la caras (*Carassius auratus gibelio* BLOCH 1782). Toxicitatea acută și subacută a fungicidului Folpan 80 WDG a fost evaluată în acvarii de sticlă. Produsul Folpan 80 WDG, în concentrațiile de la  $25 \times 10^{-5}$  Folpan g/l de apă până la  $1 \times 10^{-3}$  Folpan 80 WDG g/l de apă, produce, după o săptămână de imersiune, o scădere semnificativă a consumului de oxigen la pește. Fungicidul a schimbat ritmul respirator al peștelui în toate concentrațiile studiate, după 14 zile de expunere. Numărul de eritrocite s-a redus în mod semnificativ după 14 de zile de la imersiune, la concentrațiile fungicidului.

**Cuvinte cheie:** caras, Folpan 80 WDG, consum de oxigen, ritm respirator, număr de eritrocite.

### INTRODUCTION

Pesticides are recognized as environmental pollutants, particularly in case of ground and water (FIELDING *et al.*, 1992). The presence of pesticides has also been noted in the air by scientists and addressed by regulatory authorities (ECOBICHON, 1996). They can enter water bodies directly or indirectly, resulting in contamination of various aquatic ecosystems. The effects of insecticide pollution on non-target organisms in the environment can be studied by detecting changes in organisms at the physiological, biochemical, or molecular levels, providing “early warning” tools in monitoring environment quality (CRANE *et al.*, 1991; MIREN *et al.*, 2000). These sensitive early warning biomarkers can measure interaction between environmental xenobiotics and biological effects. Inhibition and induction of these biomarkers is a good approach to measure potential impacts of pollutants on environmental organisms (RENDO’N-VON OSTEN *et al.*, 2005).

Folpan 80WDG (active substance is folpet, (N-[(trichloromethyl)thio]phthalimide]) is a contact fungicide belonging to the dicarboximide family, a protective leaf-fungicide. Its mode of action inhibits normal cell division of a broad spectrum of microorganisms. It is used to control cherry leaf spot, rose mildew, rose black spot, and apple scab.

Folpan is a pyrethroid (SHAN *et al.*, 1997) that affects the central and peripheral nervous system and causes synaptic discharge, depolarization and ultimately death (ROBERTS & HUDSON, 1999). Pyrethroids, synthetic analogues of pyrethrins, belong to the chemical group of non-systemic insecticides (VELISEK *et al.*, 2009). Like most pyrethroids, is also an ATP-ase inhibitor. Because they are highly lipophilic, pyrethroids are likely to be strongly absorbed by the gills, even from water containing low levels of pyrethroids (SMITH & STRATTON, 1986).

Studies with typical end-use products indicate that folpet is highly toxic to both rainbow trout and bluegill sunfish. Rainbow trout were the most sensitive species and the folpet product tested was classified in the very highly toxic range of toxicity for his species. The 96-hours LC50 for bluegill sunfish is 675 ppb and the 96-hours LC50 for rainbow trout is 185 ppb (FOLPET, <http://pmep.cce.cornell.edu/profiles/extoxnet/dienochlor-glyphosate/folpet-ext.html>). Folpet is characterized as being “highly toxic” to both cold water and warm water fish.

Folpan 80 WDG is more toxic at lower temperatures, and thus more toxic to cold – than warm – water fish, but the toxicity of pyrethroids is little affected by pH or water hardness (MAUCK *et al.*, 1976).

The aim of this study is to analyze the influence of Folpan 80 WDG fungicide upon some physiological parameters (oxygen consumption, breathing frequency, number of red blood cells) in Prussian carp (*Carassius auratus gibelio*).

## MATERIAL AND METHODS

The concentrations of Folpan 80 WDG used in all experiments and treatments were established by conducting preliminary survival tests. Since the toxic substance (the folpet) is highly soluble in water, the fish individuals were introduced into water after only 5 minutes of mixing and aeration. The water temperature was maintained at 18°C-20°C, and the “immersion” solution was permanently aerated and changed every 24 hours.

Determinations were made between February and April 2011 on fish belonging to the species: Prussian carp (*C. auratus gibelio*) having an average weight of 26 g±1.28 g, 18g±0.5 g and respectively 32 g±1.4 g, caught in the surrounding lakes and rivers of Pitești city. We choose this species of fish because it is the most frequent in the Argeș River. After 10 days of adaptation in the lab, when they were fed *ad libitum* once a day, the fish were separated in lots, which were used separately for the following experiments: the first experiment was carried out with ten Prussian carp individuals having an average weight of 26 g±1.28 g, which were subjected to folpet concentrations of  $25 \cdot 10^{-5}$  Folpan 80 WDG g/l water. The second experiment was carried out with ten Prussian carp individuals having an average weight of 18g ±0.5 g, which were subjected to Folpan concentrations of  $5 \cdot 10^{-4}$  Folpan 80 WDG g/l water.

The third experiment was carried out with ten Prussian carp individuals having an average weight of respectively 32±1.4 g, which were subjected to Folpan concentrations of  $1 \cdot 10^{-3}$  Folpan 80 WDG g/l water.

The energetic metabolism, expressed by the oxygen consumption, was determined by using the closed respiratory chamber method (the oxygen dose in the water was established by using the Winkler chemical method) (PICOȘ & NĂSTĂSESCU, 1988). These determinations were made at intervals of 24, 48, 72, 96, 168 and respectively 336 hours.

The number of erythrocytes was microscopically determined with a Thoma cells numbering chamber, by using a small amount of blood collected from the caudal artery (PICOȘ & NĂSTĂSESCU, 1988).

## RESULTS

For a better comparison between the toxic effects of Folpan 80 WDG in the concentrations investigated, the average frequency of respiratory movements and oxygen consumption were represented graphically in figure 1 and figure 2.




Figure 1. The influence of Folpan 80 WDG fungicide upon oxygen consumption on Prussian carp (*Carassius auratus gibelio*). / Figura 1. Influența fungicidului Folpan 80 WDG asupra consumului de oxigen la caras (*C. auratus gibelio*).




Figure 2. The influence of Folpan 80 WDG fungicide upon breathing frequency on Prussian carp (*Carassius auratus gibelio*). / Figura 2. Influența fungicidului Folpan 80 WDG asupra frecvenței respiratorii la caras (*C. auratus gibelio*).




Figure 3. Number of erythrocytes of Prussian carp (*Carassius auratus gibelio*) after 14 days of exposure to Folpan 80 WDG fungicide. / Figura 3. Numărul eritrocitelor la caras (*C. auratus gibelio*) după 14 zile de expunere la fungicidul Folpan 80 WDG.

## DISCUSSIONS

MARINESCU *et al.*, (2004) and PONEPAL *et al.*, (2009a, 2009b) also noticed decreased oxygen consumption under the action of some pesticides and changes in respiratory rate. For all the concentrations investigated (Fig. 1), Folpan 80 WDG produced, since the first hours of exposure, a significant reduction in the oxygen consumption of the carp, as an index of their energy metabolism, reducing toxic directly proportional to the concentration and duration of exposure to its action. Folpan 80 WDG produced, in all organized experimental variants (Fig. 2), a decrease in the respiratory frequency in the case of Prussian carp, the more powerful the higher the concentration of the toxic was.

When Folpan solution was replaced every time, *C. auratus gibelio* (BLOCH 1782) was vivacious at the beginning and then relaxed. Thirty minutes later, Prussian carp became nervous, and the gills, abdomen turned red because of blood congestion; some of them were jerked, up floated and back stroked. *C. auratus gibelio* breathed quickly and their gills turned red, too because of blood congestion.

Clinical symptoms observed during Folpan exposure of Prussian carp correspond to the observations made by other authors reporting on the toxicity of pyrethroid pesticides (PRASHANTH *et al.*, 2005; VELISEK *et al.*, 2006, 2009). BRADBURY & COATS (1989) reported signs of fenvalerate poisoning in fish, which included loss of schooling behaviour, swimming near the water surface, hyperactivity, erratic swimming, seizures, loss of buoyancy, increased cough rate, increased gill mucus secretions, flaring of the gill arches, head shaking and listlessness before death. Sublethal effects of pyrethroids on fish include gill damage and behavioural changes (VELISEK *et al.*, 2009). In fish, direct contact between the aquatic environment and the gill epithelium may cause these surfaces to become sensitive to environmental alteration in the presence of toxic materials or other irritants.

Folpan effects on Prussian carp are positively related to the concentrations of the chemical substances, suggesting an obvious dose-response relationship. As for the mortalities of  $25 \cdot 10^{-5}$ g Folpan 80 WDG, only one piece of Prussian carp died on the 10<sup>th</sup> day. In  $5 \cdot 10^{-4}$ g Folpan 80 WDG concentration the mortality of Prussian carp was 1 on the 12<sup>th</sup> day and increased to 3 on the 14<sup>th</sup> day. The mortality of  $1 \cdot 10^{-3}$ g Folpan 80 WDG was 100% on the 14<sup>th</sup> day. The experimental results showed that mortality was very small in the case of lower concentrations of Folpan 80 WDG, and was raised significantly as the concentrations of Folpan 80 WDG increased; when the concentrations of Folpan 80 WDG reached a higher value, all Prussian carp individuals were seriously toxic and died in a very short period of time. It is suggested that water organisms can be poisoned by Folpan 80 WDG such as Prussian carp, and toxicity relies on the chemical concentrations and the exposure duration.

The toxic effect of Folpan was proven to be more powerful in the first 24 hours from the fish immersion.

Respiratory irregularities are thought to be caused by mucus precipitation on the gill epithelium in response to a toxicant (SCHAUMBURG *et al.*, 1967). The use of respiratory stress to monitor sublethal effects of intoxication was previously applied to a variety of toxicants and subjects (SCHAUMBURG *et al.*, 1967; WALDEN *et al.*, 1970). This may result in a decrease in the dissolved oxygen at the gill surface, initiating the cough reflex which is an attempt to clean the respiratory surface.

Haematological studies in fish have assumed greater significance because these parameters were to be used as an effective and sensitive index to monitor physiological and pathological changes induced by natural or anthropometric factors (DHEMBARE *et al.*, 2000). Prussian carp anaemia could be due to hypoxia that was induced by injuring the gills, as the red-pink colour of the gills became red-white, and at high concentrations the gills completely lost their red colour, while abundant secretions of mucus and even mucosal detachment with abundant bleeding could be observed (Fig. 3).

## CONCLUSIONS

Folpan 80 WDG produced, in all organized experimental variants a decrease in respiratory frequency and consumption of oxygen in the case of Prussian carp, the concentration is higher as the more powerful of the toxic was.

In all experimental variants that have been applied on fish, all the stages described by Schäperclaus in the symptomatology scheme for the intoxicated fish (PICOȘ & NĂSTĂSESCU, 1988), were only observed in the variant with the Folpan 80 WDG in concentration of  $1 \cdot 10^{-3}$ g/l water, where they succeeded each other at very short intervals (the fish died in the first 24 experimental hours). For the other variants we only noticed the first three stages.

After two weeks of exposure to  $5 \cdot 10^{-4}$ g Folpan 80 WDG the number of erythrocytes in Prussian carp decreased significantly compared to the control groups.

## ACKNOWLEDGMENT

This work was partially supported by the strategic grant POSDRU 107/1.5/S/77265 (2010) of the Ministry of Labour, Family and Social Protection, Romania, co-financed by the European Social Fund – Investing in people.

## REFERENCES

- BRADBURY S. P. & COATS J. R. 1989. *Comparative toxicology of the pyrethroid insecticides*. Bulletin of Environmental Contamination and Toxicology. SpringerLink. **108**: 134-177.

- CRANE M. & MALTBY LORRAINE. 1991. *The lethal and sub-lethal responses of *Grammarus pulex* to stress: sensitivity and sources of variation in a situ bioassay*. Environmental Toxicology and Chemistry. Wiley-Blackwell. **10**: 1331-1339.
- DHEMBARE A. J. & PONDHA G. M. 2000. *Haematological changes in fish *Puntius sophore* exposed to some insecticide*. Journal of Experimental Zoology. Elsevier. India. **3**(1): 41-44.
- ECOBICHON D. J. 1996. *Toxic effects of pesticides*. In: Klaasen CD. (Eds.) Casarett and Doull's Toxicology: The Basic Science of Poisons. 5<sup>th</sup> Edit. New York: McGraw Hill. 1310 pp.
- FIELDING M., BARCELO D., HELWEG A., GALASSI S., TORSTENSSON L., VAN ZOONER P., WOLTER R., ANGELETTI G. 1992. *Pesticides in ground and drinking water*. Edit. Water Pollution Research Report 27. In: Commission of the European Communities Brussels. 1136 pp.
- MARINESCU AL. G., DRĂGHICI O., PONEPAL CRISTINA, PĂUNESCU ALINA. 2004. *The influence of fungicide (Dithane M-45) on some physiological indices in the prussian carp (*Carassius auratus gibelio* Bloch)*. International Association for Danube Research. Novi Sad. **35**: 209-214.
- MAUCK W. L., OLSON L. E., MARKING L. L. 1976. *Toxicity of natural pyrethrins and five pyrethroids to fish*. Archives of Environmental Contamination and Toxicology. Springer Link. **4**: 18-29.
- MIREN P., CAJARAVILLE M. J., BEBIANNO J. B., CINTA P., CARMEN S., ALDO V. 2000. *The use of biomarkers to assess the impact of pollution in coastal environments of the Iberian Peninsula: a practical approach*. Science of the Total Environment. Mendeley. **247**: 295-311.
- PICOȘ C. A. & NĂSTĂSESCU GH. 1988. *Lucrări practice de fiziologie animală*. Edit. Universității din București. 232 pp.
- PONEPAL MARIA CRISTINA, PĂUNESCU ALINA, MARINESCU AL. G., DRĂGHICI O. 2009a. *The Changes of Some Physiological Parameters in Prussian Carp Under The Action of the Tilt Fungicide*. Bulletin USAMV. Cluj-Napoca. **66**(1-2): 47-52.
- PONEPAL MARIA CRISTINA, PĂUNESCU ALINA, MARINESCU AL. G., DRĂGHICI, O. 2009b. *Effect of the Fungicide Chlorothalonil (Bravo) on Some Physiological Parameters in Prussian Carp*. Lucrări științifice USAMV Iași. seria Horticultură. **52**: 1157-1162.
- PRASHANTH M. S., DAVID M., MATHED S. G. 2005. *Behavioral changes in freshwater fish, *Cirrhinus mrigala* (Hamilton) exposed to cypermethrin*. Journal of Environmental Biology. Academy of Environmental Biology India. Mendeley. **26**: 141-144.
- RENDO'N-VON OSTEN J., ORTIZ-ARANA A., GUILHERMINO L., SOARES A. M. 2005. *In vivo evaluation of three biomarkers in the mosquitofish (*Gambusia yucatana*) exposed to pesticides*. Chemosphere. Elsevier. **58**: 627-636.
- ROBERTS T. & HUDSON D. 1999. *Metabolic pathway of agrochemicals*. Part 2: insecticides and fungicides. 1<sup>st</sup> Edit. The Royal Society of Chemistry. Cambridge. United Kingdom. 849 pp.
- SCHAUMBURG F. D., HOWARD T. E., WALDEN C. C. 1967. *A method to evaluate the effects of water pollution on fish respiration*. Water Research. Science Direct. **1**: 731-737.
- SHAN G., HAMMER R. P., OTTEA J. A. 1997. *Biological activity of pyrethroid analogs in pyrethroid-susceptible and-resistant tobacco budworms, *Heliothis virescens* (F.)*. Journal of Agricultural and Food Chemistry. Agris. **45**: 4466-4473.
- SMITH T. M. & STRATTON G. W. 1986. *Effects of synthetic pyrethroid insecticides on nontarget organisms*. Research Reviews. Mendeley. **97**: 93-119.
- VELISEK J., DOBSIKOVA R., SVOBODOVA Z., MODRA H., LUSKOVA V. 2006. *Effect of deltamethrin on the biochemical profile of common carp (*Cyprinus carpio* L.)*. Bulletin of Environmental Contamination and Toxicology. Mendeley. **76**: 992-998.
- VELISEK G., SVOBODOVA Z., MACHOVA J. 2009. *Effects of bifenthrin on some haematological, biochemical and histopathological parameters of common carp (*Cyprinus carpio* L.)*. Fish Physiology and Biochemistry. Springer Link. **35**: 583-590.
- WALDEN C. C., HOWARD T.E., FROUD G.C. 1970. *A quantitative assay of the minimum concentration of kraft mill effluents which affect fish respiration*. Water Research. Elsevier. **4**: 61-68.
- \*\*\*. *A Pesticide Information Project of Cooperative Extension Offices of Cornell University*. Oregon State University. The University of Idaho And the University of California at Davis and the Institute for Environmental Toxicology. Michigan State University. *Folpet*. <http://pmep.cce.cornell.edu/profiles/extoxnet/dienochlor-glyphosate/folpet-ext.html> (accessed in March 3, 2012).

Zgurschi Gabriela, Păunescu Alina, Marinescu Al. Gabriel

University of Pitești, Târgu din Vale Street, nr.1, 410087,

Pitești, Romania

E-mail: gabrielazgurschi@yahoo.com

Received: March 31, 2012

Accepted: June 20, 2012