

ARE HUMAN SETTLEMENTS ECOLOGICAL SYSTEMS?

PETRIȘOR Alexandru-Ionuț

Abstract. Even though “human ecology” is an old and often used concept, many debates are present in the ecological literature whether human settlements are ecological systems or not. The paper analyses the evolution, structure and functions of human settlements in relationship to their scale. The results indicate that, despite of the altered structure and functions resulting from the presence and activities of the dominant human species, human settlements are ecological systems, ranging based on their size from “ecosystems” to “complexes of ecosystems” (landscapes).

Keywords: urban ecosystem, functional structure, biodiversity, human species, scale.

Rezumat. Sunt așezările umane sisteme ecologice? Deși termenul „ecologie urbană” datează de ceva vreme și este des folosit, în literatura de specialitate din domeniul ecologiei sunt prezente multe dezbateri privind posibilitatea de a considera așezările umane sisteme ecologice. Lucrarea analizează evoluția, structura și funcțiile așezărilor umane în funcție de scara acestora. Rezultatele arată că, în pofida modificării structurii și funcțiilor datorită prezenței și activității speciei umane dominante, așezările umane sunt sisteme ecologice, situate în funcție de mărime pe nivelul ecosistemelor sau complexelor de ecosisteme.

Cuvinte cheie: ecosistem urban, structură funcțională, biodiversitate, specia umană, scară.

INTRODUCTION

Analysing different definitions of “systems”, it can easily be seen that the concept describes a sum of elements working together as a whole (BOTNARIUC & VĂDINEANU, 1982); the elements are objects and their relationships (IANOȘ & HELLER, 2006). Other authors consider that systems are functional structures (IANOȘ, 2000; VĂDINEANU, 1998, 2004; PETRIȘOR, 2008). The later view expands the first one showing that the structure (interrelated elements) is adapted to its functions in a double sense: functions modify the structure, and a certain structure can carry only one or more specific functions (PETRIȘOR, 2011). This feature characterizes mechanical systems (e.g., a car must have a certain structure in order to function as a vehicle), living systems and other systems too. In ecology, Tansley’s definition (TANSLEY, 1935) identifies the two components of an ecosystem: the living component (called in the British literature “biocoenosis” and in the American one “community”) and the non-living component (biotope).

While “urban ecology” is said to trace its roots in the 50’s, but its first dated use was during the 1968 symposium “Challenge for survival in megalopolis” (PETRIȘOR, 2008). Since ecology embraced the systemic theory, urban ecology followed the same pattern and authors started using the term “urban ecosystem”. In 1997, a new journal, “Urban Ecosystems”, was started in the Netherlands. The first Editorial defines the urban ecosystem as a particular type of ecosystem (based on Tansley’s view), where man is the “*keystone species controlling ecosystem structure and function*” (WALBRIDGE, 1997).

The dominant presence of humans in their settlements, and the control exercised over the structure and functions of ecosystems made many authors reluctant in accepting the fact that human settlements are ecosystems (METZGER, 1994; MCINTYRE et al., 2000; REES, 2003; PICKETT & GROVE, 2009; COLLINS et al., 2012), while others seem to embrace the new concept (CRISTEA & BACIU, 2000; EUROPEAN ENVIRONMENT AGENCY, 2010). One of the underlying causes could be the lack of interest for carrying out research in inhabited areas (MARRIS, 2009; CORBYN, 2010; COLLINS et al., 2012). Other authors underline structural differences, particularly related to biodiversity (CRISTEA & BACIU, 2000; SAVARD et al., 2000; EUROPEAN ENVIRONMENT AGENCY, 2010; ȘUSTEK, 2011, 2012; RIDICHE & BĂLESCU, 2012) or altered biotope conditions due to human activities (GAVRILESCU, 2011; CORNEANU et al., 2012), functional differences (BOLUND & HUNHAMMAR, 1999; CRISTEA & BACIU, 2000; DECKER et al., 2000; LUNDHOLM, 2006), or scale (METZGER, 1994; SAVARD et al., 2000; CLERGEAU et al., 2006).

Especially in regard to the scale, based on Gaia theory (LOVELOCK, 1979), other authors assimilate the city, despite of its size, with an organism, and discuss about the “urban metabolism” (DECKER et al., 2000; CRĂCIUN, 2008; GOLUBIEWSKI, 2012), up to identifying “normal” and “pathological” forms of it (STAN, 2011). At the opposite pole, other authors consider that cities are situated at the level of “complexes of ecosystems” or “landscapes” (METZGER, 1994; CLERGEAU et al., 2006; PETRIȘOR & PETRIȘOR, 2008; PETRIȘOR, 2010; 2011).

The aim of this study is to discuss the structure and functions of human settlements from an ecological perspective, in order to see whether they can be assimilated to ecological systems.

EVOLUTION

During the historical evolution, human settlements were first similar to what we call “rural” nowadays. Gradually, some of them became “urban areas”. Therefore, the anthropization process consists of a first transformation of natural systems into rural systems, and later into urban systems. The presence of humans is more intense in cities,

resulting into additional elements that exacerbate the difference between natural and man-dominated systems along the gradient natural – rural – urban; for this reason, the conceptual model displayed in figure 1 shows a transversal line symbolizing the balance between natural and anthropic elements, leaning towards the latest in urban systems.

In this context, it is noteworthy mentioning that the definition of “urban” areas has a different meaning for ecologists, economists, sociologists, psychologists or planners (MCINTYRE et al., 2000); in general, the United States definitions are based on the density of human population, while the European ones focus on key indicators (such as educational or cultural infrastructure, but also from other social or economic areas) (PETRIȘOR, 2008).

During the process, man-dominated systems couple functionally to the natural ones. Resources are taken directly or from agro-ecosystems, a component of the socio-economic system, using technology. Energy is present in fertilizers, pesticides, soil and green space works. Humans change the structure of geographic spaces. Altered biogeochemistry and loss of biodiversity decrease the stability of natural systems and increased their dependence on the man-dominated ones (COLLINS et al., 2012). The process is assessed using the concept of eco-energy – initial energy, before conscious human interventions (IANOȘ, 2000); the concentration of population and economic activities consume primary eco-energies as the level of anthropization and complexity of geosystems increase. Eco-diversity and geodiversity increase at the expense of biodiversity (IANOȘ et al., 2011; PETRIȘOR & SÂRBU, 2010).

STRUCTURE

Even though the structure of natural, rural and urban systems does not differ in terms of the names of components, detailed variation is visible along gradients of anthropization (ȘUSTEK, 2011; 2012). In the introductory section, the living component of an ecosystem was called in the British literature “biocoenosis” and in the American one “community”. The latter term can produce confusions, especially when authors discuss about “mammal communities” or “bird communities”. The biocoenosis consists of all plant and animal species of an ecosystem (BOTNARIUC & VĂDINEANU, 1982); “community” can be used equivalently or in order to refer to a specific group. Especially in man-dominated systems, fragmentation, loss of biodiversity and functional disturbances induced by human activities determine the assembly of species to work as a group of “communities” instead of being a whole.

The overall biodiversity of species is lower than in natural systems, resulting into shorter food webs, and into the fact that man-dominated systems are “incomplete” and the reduced biodiversity prevents the specialization of its components (PETRIȘOR, 2008). However, some groups (particularly invertebrates) are favoured. Man replaces species in time introducing new ones or eliminating the existing ones directly or indirectly; as a result, the entire biocoenosis is changed over longer periods. Researchers have found a variety of niches and habitats even in urban areas (PETRIȘOR, 2008; 2010). They are inhabited by species seeking for human habitats – so-called hemerophilous species, including the ones that are found only in these areas (synanthropic), indifferent to the presence of humans (hemerodiaphore) (NOBLET, 1994, 2005; PETRIȘOR, 2008, 2010). From a different viewpoint, man-dominated systems include many ubiquitous and opportunistic species (the latter can be eventually favoured by shifting conditions due to human activities), and to a lesser extent by random species (CLERGEAU et al., 2006; PETRIȘOR, 2008, 2010). In addition to them, people bring allochthonous species, some even invasive (PETRIȘOR, 2008; COLLINS et al., 2012), and proliferate the domestic species (PETRIȘOR, 2008). Most often, the species characterizing natural systems from the same region are confined to the green spaces (PETRIȘOR, 2008; EUROPEAN ENVIRONMENT AGENCY, 2010). In a spatial perspective, the type of biodiversity (α , β etc.) can be correlated with the size of the human settlement and described in relationship with the diversity of habitats, land use or cover, biogeographical region, etc. (PETRIȘOR, 2012a). However, if biodiversity is diminished, the diversity of man-generated structures is increased with the level of anthropization.

The biotope is altered too. Over long periods of time, even the geography is changed by modifying water courses and through the pressure exercised by the increased mass of constructions. The soil is removed or covered by asphalt, or, if it exists, does not have too much to offer to vegetation (PETRIȘOR, 2008). Microclimate is affected especially by urban areas, which become heat islands, particularly during the warm season (CHEVAL et al., 2009; EUROPEAN ENVIRONMENT AGENCY, 2010; COLLINS et al., 2012) due to the alteration of biogeochemical cycles, such as the water circuit discussed in the next section. However, these changes are the least evident. What becomes obvious in man-dominated system is the presence of infrastructure (built capital, physical capital) and pollutants (METZGER, 1994; REES, 2003; PETRIȘOR, 2008, 2010; PICKETT & GROVE, 2009; GAVRILESCU, 2011; CORNEANU et al., 2012). All these are influenced by what could be called the “human mind”, and assessed by psychological, social, anthropological, cultural structures, resulting into the socio-economic, political, legislative and administrative conditions that set their fingerprint over the layout and operation of human settlements (BOȘTENARU, 2005; PETRIȘOR, 2008, 2010; PICKETT & GROVE, 2009).

The separation between man-dominated systems and the adjacent ones is naturally done by ecotone areas. Researchers have identified urban fringes as having the same function (STAN, 2009; COLLINS et al., 2012). However, based on its social analysis, other authors consider that urban fringes have a separating role instead of a joining one, and certainly do not account for the productivity of the two ecosystems (PETRIȘOR, 2012b).

Figure 1 displays the components of human settlements in this view, as well as their connections. Based on the definition of territorial systems (IANOȘ, 2000) and ecological standpoint (TANSLEY, 1935; BOTNARIUC & VĂDINEANU, 1982; VĂDINEANU, 2004; PETRIȘOR, 2008, 2010, 2011), the figure has four areas, corresponding to the division between “natural” vs. “man-dominated” and “living” vs. “non-living”.

FUNCTIONS

Again, in theory man-dominated systems carry out the same functions as the natural ones: biogeochemical cycles and self-regulation. Nevertheless, they are radically changed. Biogeochemical cycles are disturbed, often resumed to a straight, unidirectional linear flow (CRISTEA & BACIU, 2000; PETRIȘOR, 2008). Some of the examples relate to the water circuit: in cities, water reaches the sewerage system, and is taken outside; consequently, urban planners find a strong need to create artificial water bodies or courses, or at least fountains (CRISTEA & BACIU, 2000; DECKER et al., 2000). Also, the input of natural energy is diminished due to the scarcity of primary producers. Consequently, man-dominated systems need external matter and energy sources, found in the natural ones (CRISTEA & BACIU, 2000; DECKER et al., 2000). From this perspective, they become “energetic parasites” of these systems (VĂDINEANU, 2004; PETRIȘOR, 2008). The dissipative character is more evident in man-dominated system than in the natural ones, due to the structuring interventions of humans (REES, 2003; PETRIȘOR & SĂRBU, 2010).

Moreover, the food levels can be seen as part of an entire hierarchy of levels, starting with the traditional trophic ones, derived from physical consumption of food, and continuing with technotrophy (consumption of energy and resources by technology) and nootrophy (their consumption in the activities supporting human development, such as research, education, management, administration etc.) (IANOȘ, 2000; REES, 2003; PETRIȘOR, 2011).

Self-regulation is inhibited by the loss of biodiversity (CRISTEA & BACIU, 2000; EUROPEAN ENVIRONMENT AGENCY, 2010) and, in general, by the control exercised by humans. If the evolution of general systems is described by concepts like “succession” (BOTNARIUC & VĂDINEANU, 1982; DECKER et al., 2000; MCINTYRE et al., 2000), “panarchy” or “adaptive cycles” (HOLLING, 2004; VĂDINEANU, 2004), the model proposed for the human settlements is a spiral, suggesting the fact that humans change the characteristics of their system through each intervention (IANOȘ et al., 2011; PETRIȘOR, 2011). Nevertheless, from the succession standpoint, man-dominated systems are young, immature (PETRIȘOR & IANOȘ, 2011).

Functions are also affected by the action of natural laws. One of the limiting ones is the “minimal law”, meaning that many parameters required for the normal life of species are reduced to a minimum and act as constraining factors. Mitscherlich’s law, describing the decrease in intensity of the favourable action of some factors as their dosage increases, explains the need to use extra-energy in order to maintain natural areas like urban green spaces (PETRIȘOR, 2008).




Figure 1. Conceptual model of the structure of man-dominated systems in relationship to their functions.

Last but not least, specific functions oriented by the role of human settlements (military, commercial, capital etc.) influence their structure and functions seen from an ecological perspective, resulting into diminishing or favouring some of them, but also in increased pollution (PETRIȘOR, 2008; PEPTENATU et al., 2010, 2012), depending on the attitude towards the environment (IANOȘ et al., 2009).

Figure 1 displays the functional characteristics of human settlements, with a particular focus on energy. The focus is determined by the relationship between energy, land use, climate change and composition of biocoenoses, and its consequences over the planning process (DALE, 1997; BOLUND & HUNHAMMAR, 1999; DALE et al., 2009, 2011; MEIȚĂ et al., 2011; PETRIȘOR et al., 2011; PETRIȘOR, 2012a). Land cover refers to a biophysical interpretation of what lies on the ground surface (JENSEN, 2000), while land use indicates its use by human communities or a more detailed classification of natural systems (PETRIȘOR et al., 2010).

SCALE

Apart from the diversity of habitats, human settlements are, from the standpoint of land cover and use, a mosaïque of vegetated areas (pastures within rural areas, green spaces within the urban ones), semi-natural areas (parks, green spaces, encroached portions of natural systems), human-built infrastructure and occasionally other elements. For this purpose, some authors suggest that, on a scale starting with ecosystems, continuing with the complex of ecosystems or landscapes and ending with the planetary system – ecosphere, large human settlements, especially cities, through their complexity exceed the first level approaching the second (METZGER, 1994; BOLUND & HUNHAMMAR, 1999; CLERGEAU et al., 2006; PETRIȘOR, 2010, 2011).

At the next spatial level, man-dominated system and adjacent infrastructure (roads) form the socio-economic system, expanding over the natural ones. The characteristics of its dynamics are: (1) spatial expansion by substituting, simplifying and fragmenting natural systems; (2) increase of inner complexity and fluxes of resources taken from natural systems and pollutants dispersed into them; (3) linearization of biogeochemical cycles; (4) accumulation of waste; and (5) regionalization and globalization through increased connectivity (VĂDINEANU, 1998).

CONCLUSIONS

The paper aimed to answer the question whether man dominated systems, particularly human settlements, can still be considered ecological systems. Many differences have been found between the extremes of an evolution scale starting with natural system, including the rural systems and ending with the urban systems, in order to reflect the degree of anthropization and consumption of natural energy and resources, but also between the three levels. The differences are found in their structure (loss of biodiversity and presence of man-induced infrastructure and pollutants in man-dominated systems) and functions (linearization of biogeochemical cycles, increased dependence of human activities, energy and resources of other systems, and low self-regulation capacity of man-dominated systems). Despite of these differences, the essential components of ecological systems are still present. Based on this, human settlements can be considered ecological systems with a different homomorphous model, situated, depending on their scale, at the level of “ecosystems” or “complexes of ecosystems” (landscapes).

ACKNOWLEDGEMENT

Thanks are due to Professors Mircea VARVARA and Ștefan NEGREA for their valuable comments and suggestions, which have contributed to improving the quality of the manuscript.

REFERENCES

- BOLUND P. & HUNHAMMAR S. 1999. *Ecosystem services in urban areas*. Ecological Economics. Amsterdam. **29**: 293-301.
- BOȘTENARU D. M. 2005. *Multidisciplinary co-operation in building design according to urbanistic zoning and seismic microzonation*. Natural Hazards and Earth System Sciences. Cuvillier Verlag. Göttingen. **5**: 397-411.
- BOTNARIUC N. & VĂDINEANU A. 1982. *Ecologie*. Edit. Didactică și Pedagogică. București. 438 pp.
- CHEVAL S., DUMITRESCU A., BELL A. 2009. *The urban heat island of Bucharest during the extreme high temperatures of July 2007*. Theoretical and Applied Climatology. Springer-Verlag. Heidelberg. **97**(3-4): 391-401.
- CLERGEAU P., JOKIMÄKI J., SNEP R. 2006. *Using hierarchical levels for urban ecology*. Trends in Ecology and Evolution. Maryland Heights. MO. **21**(12): 660-661.
- COLLINS J. P., KINZIG A., GRIMM N. B., FAGAN W. F., HOPE D., WU J., BORER E. T. 2012. *A New Urban Ecology. Modeling human communities as integral parts of ecosystems poses special problems for the development and testing of ecological theory*. American Scientist. Research Triangle Park. NC. **88**: 416-425.
- CORBYN Z. 2010. *Ecologists shun the urban jungle*. Nature. London. doi:10.1038/news.2010.359.
- CORNEANU M., CORNEANU G., COJOCARU L., LĂCĂTUȘU A. R. 2012. *Investigations to detect ecosystem disturbances under the influence of anthropogenic factors*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **28**(2): 219-228.
- CRĂCIUN C. 2008. *Metabolismul urban. O abordare neconvențională a organismului urban*. Edit. Universitară „Ion Mincu”. București. 366 pp.
- CRISTEA V. & BACIU C. 2000. *Une approche écosystémique de la ville*. Naturopa. Bietlot-Gilly. **94**: 4-5.

- DALE V. H. 1997. *The relationship between land-use change and climate change*. Ecological Applications. ESA Publications Office. Ithaca. NY 7(3): 753-769.
- DALE V. H., EFROYMNSON R. A., KLINE K. L. 2011. *The land use–climate change–energy nexus*. Landscape Ecology. Springer. New York. 26: 755-773.
- DALE V. H., LANNOM K. O., THARP M. L., HODGES D. G., FOGEL J. 2009. *Effects of climate change, land-use change, and invasive species on the ecology of the Cumberland forests*. Canadian Journal of Forest Research. Research Press. Montreal. 39: 467-480.
- DECKER E. H., ELLIOTT S., SMITH F. A., BLAKE D. R., ROWLAND F. S. 2000. *Energy and material flow through the urban ecosystem*. Annual Review of Environment and Resources. Published Heinz et Miniwarter. Palo Alto. 25: 685-740.
- EUROPEAN ENVIRONMENT AGENCY. 2010. *10 messages for 2010. Urban ecosystems*. EEA. Copenhagen. 11 pp.
- GAVRILESCU ELENA. 2011. *The study of air quality indices in Craiova Municipality*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. 27(1): 155-158.
- GOLUBIEWSKI N. 2012. *Is There a Metabolism of an Urban Ecosystem? An Ecological Critique*. Ambio. Heidelberg. 2012: 1-14.
- HOLLING C. S. 2004. *From complex regions to complex worlds*. Ecology and Society. Wolfville. Nova Scotia. Canada 9(1). On-line at <http://www.ecologyandsociety.org/vol9/iss1/art11/> (accessed January 7, 2013)
- IANOȘ I. 2000. *Sisteme teritoriale. O abordare geografică*. Edit. Tehnică. București. 197 pp.
- IANOȘ I. & HELLER W. 2006. *Spațiu, economie și sisteme de așezări*. Edit. Tehnică. București. 373 pp.
- IANOȘ I., PEPTENATU D., ZAMFIR D. 2009. *Respect for environment and sustainable development*. Carpathian Journal of Earth and Environmental Sciences. Edit. Universității Baia Mare. 4(1): 81-93.
- IANOȘ I., PETRIȘOR A.-I., STOICA I. V., SÂRBU C. N., ZAMFIR D., CERCLEUX A. L. 2011. *The different consuming of primary eco-energies and their degradation in territorial systems*. Carpathian Journal of Earth and Environmental Sciences. Edit. Universității Baia Mare. 6(2): 251-260.
- JENSEN J. R. 2000. *Remote Sensing of the Environment. An Earth Resource Perspective*. Prentice Hall. Upper Saddle River. NJ. 544 pp.
- LOVELOCK J. E. 1979. *Gaia: A new look at life on Earth*. Oxford University Press. Oxford. 157 pp.
- LUNDHOLM J. T. 2006. *How novel are urban ecosystems?* Trends in Ecology and Evolution. Maryland Heights. 21(12): 659-660.
- MARRIS E. 2009. *Ragamuffin Earth*. Nature. London University Press. London. 460: 450-453.
- MCINTYRE N. E., KNOWLES-YÁNEZ K., HOPE D. 2000. *Urban ecology as an interdisciplinary field: differences in the use of "urban" between the social and natural sciences*. Urban Ecosystems. Publisher Springer. Heidelberg. 4: 5-24.
- MEIȚĂ V., PETRIȘOR A.-I., SIMION-MELINTE C.-P. 2011. *Agricultural impact of the exposure to climate change in the Romanian portion of Tisza river basin*. Research Journal of Agricultural Science. Publishing House EUROBIT. Timișoara. 43(3): 429-436.
- METZGER G. 1994. *Contribution à une problématique de l'environnement urbain*. Cahiers des sciences humaines. Annales de la Faculté des lettres et des sciences humaines. Paris. 30(4): 595-619.
- NOBLET J.-F. 1994. *La maison nichoir. Hommes et bêtes: comment cohabiter?* Edit. Terre vivante. Paris. 128 pp.
- NOBLET J.-F. 2005. *La Nature sous son toit. Hommes et bêtes: comment cohabiter?* Delachaux et Niestlé. Paris. 173 pp.
- PEPTENATU D., PINTILII R. D., DRAGHICI C., STOIAN D. 2010. *Environmental pollution in functionally restructured urban areas: case study – the city of Bucharest*. Iranian Journal of Environmental Health Science & Engineering. Published by Tehran University. Tehran. 7(1): 87-96.
- PEPTENATU D., MERCIU C., MERCIU G., DRĂGHICI C., CERCLEUX L. A. 2012. *Specific Features of Environment Risk Management in Emerging Territorial Structures*. Carpathian Journal of Earth and Environmental Sciences. Edit. Universității Baia Mare. 7(2): 135-143.
- PETRIȘOR A. I., IANOȘ I., TĂLÂNGĂ C. 2010. *Land cover and use changes focused on the urbanization processes in Romania*. Environmental Engineering and Management Journal. Edit. Universității Iași. 9(6): 765-771.
- PETRIȘOR A. I. & IANOȘ I. 2011. *Application of the methods used in the study of biological diversity to territorial systems*. Analele Universității Ovidius. Seria Geografie. Constanța. 5(1): 27-34.
- PETRIȘOR A. I. & PETRIȘOR L. E. 2008. *Integrarea în peisaj, UICN și Convenția de la Florența*. logiA. Cluj Napoca. 10: 113-117.
- PETRIȘOR A. I., MEIȚĂ V., CHICOȘ A., PELEANU I., SIMION-MELINTE C.-P. 2011. *Assessing the vulnerability to climate change in the Romanian part of Tisza river basin*. Romanian Review of Regional Studies. Edit. Universității Cluj Napoca. 7(2): 121-128.
- PETRIȘOR A. I. & SÂRBU C. N. 2010. *Dynamics of geodiversity and eco-diversity in territorial systems*. Journal of Urban and Regional Analysis. Publisher University of Bucharest. 2(1): 61-70.
- PETRIȘOR A. I. 2008. *Ecologie urbană, dezvoltare spațială durabilă și legislație*. Edit. Fundației România de mâine. București. 272 pp.
- PETRIȘOR A. I. 2010. *Mediul urban: o abordare ecologică*. Urbanistique. On-line at <http://www.urbanistique.ro/mediul-urban-o-abordare-ecologica-dr-alexandru-ionut-petrisor/> (Accessed: January 7, 2013).

- PETRIȘOR A. I. 2011. *Systemic theory applied to ecology, geography and spatial planning. Theoretical and methodological developments*. Lambert Academic Publishing. Saarbrücken. 172 pp.
- PETRIȘOR A. I. 2012a. *Land cover and use changes and predicted climate changes in Romania: Connections underlined by their spatial distributions*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **28**(1): 141-148.
- PETRIȘOR A. I. 2012b. *Land cover and land use analysis of urban growth in Romania*. Human Geographies. Publisher University of Bucharest. **6**(1): 47-51.
- PICKETT S. T. A. & GROVE J. M. 2009. *Urban ecosystems: What would Tansley do?* Urban Ecosystems. Publisher Springer. Heidelberg. **12**: 1-8.
- REES W. E. 2003. *Understanding Urban Ecosystems: An Ecological Economics Perspective*. In: Berkowitz A., Nilon C., Hollweg K. Understanding Urban Ecosystems. Springer-Verlag. New York: 115-136.
- RIDICHE MIRELA SABINA & BĂLESCU CARMEN DANIELA. 2012. *Preliminary study on the avifauna in Radovan locality area (Dolj County, Romania)*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **28**(2): 123-132.
- SAVARD J.-P. L., CLERGEAU P., MENNECHEZ G. 2000. *Biodiversity concepts and urban ecosystems*. Landscape and Urban Planning. Amsterdam. **48**: 131-142.
- STAN A. I. 2009. *Peisajul periferiilor urbane: revitalizarea peisageră a zonelor periferice*. Edit. Universitară „Ion Mincu”. București. 295 pp.
- STAN A. I. 2011. *Maladii urbane in perioada crizei economice*. Urbanism. Arhitectură. Construcții. Edit. Astra. Craiova. **3**(3): 17-24.
- ŠUSTEK Z. 2011. *Changes in carabid communities (Insecta: Coleoptera) along an urbanization gradient in Pyongyang (North Korea)*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **27**(1): 73-92.
- ŠUSTEK Z. 2012. *Changes in carabid communities along the urbanization gradient in Madrid (Spain)*. Oltenia. Studii și comunicări. Științele Naturii. Muzeul Olteniei Craiova. **28**(2): 87-96.
- TANSLEY A. G. 1935. *The use and abuse of vegetational concepts and terms*. Ecology. ESA Publications Office. Ithaca. NY. **16**(2): 284-307.
- VĂDINEANU A. 1998. *Dezvoltarea durabilă. Bazele teoretice ale dezvoltării durabile*. Edit. Universității din București. **1**. 248 pp.
- VĂDINEANU A. 2004. *Managementul dezvoltării: o abordare ecosistemică*. Edit. Ars Docendi. București. 394 pp.
- WALBRIDGE M. R. 1997. *Urban Ecosystems*. Urban Ecosystems. Publisher Springer. Heidelberg. **1**: 1-2.

Petrișor Alexandru-Ionuț

Department of Urban and Landscape Planning, School of Urbanism and Landscape Architecture

“Ion Mincu” University of Architecture and Urbanism, Bucharest

Str. Academiei. no. 18-20. sector 1. cod 010014. Bucharest. Romania.

National Institute for Research and Development in Constructions, Urbanism and Sustainable Spatial Development URBAN-INCERC

Șos. Pantelimon no. 266. sector 2. cod 021652. Bucharest. Romania.

E-mail: alexandru_petrisor@yahoo.com, Internet: www.environmentics.ro

Received: January 10, 2013

Accepted: May 8, 2013