

COMPARATIVE DATA ON PALEO-BIODIVERSITY OF GASTROPODS IN LATTAKIA REGION, SYRIA IN CORELATION WITH THEIR LIVING ENVIRONMENTS

AXINI Monica, AL-AZKI Fawaz

Abstract. Lattakia is a region located on the eastern coast of the Mediterranean Sea. It is a great and important area from a geological, geomorphological, biogeographical, palaeontological and biodiversity point of view, in many scientific and practical aspects. This paper presents data on the diversity of fossil gastropods identified so far in Lattakia Region in correlation with the current situation. Also, this paper is intended to be a reconstruction of palaeo-environment during the Cretaceous in Lattakia Region today. These data represent preliminary results of research carried out during 1985-2014 in Region Lattakia, Syria.

Keywords: fossil gastropods, palaeo-biodiversity, data of palaeo-ecology, Lattakia (Syria).

Rezumat. Date comparative privind paleo-biodiversitatea gasteropodelor din Regiunea Lattakia, Siria în comparație cu mediile lor de viață. Lattakia este o regiune situată pe coasta estică a Mării Mediterane. Este o zonă deosebită din punct de vedere geologic, geomorfologic, biogeografic, paleontologic și al biodiversității, și importantă prin multe aspecte științifice și practice. Lucrarea prezintă date privind diversitatea gasteropodelor fosile identificate până în prezent din Regiunea Lattakia în corelație cu situația actuală. Lucrarea se dorește a fi și o reconstituire a paleomedului din timpul Cretacului în regiunea Lattakiei de astăzi. Aceste date reprezintă rezultate preliminare ale cercetărilor desfășurate în perioada 1985-2014 în Regiunea Lattakia, Siria.

Cuvinte cheie: gasteropode fosile, paleo-biodiversitate, date de paleo-ecologie, Lattakia (Syria).

INTRODUCTION

Lattakia Region is located in northwest Syria, in the Middle East of Asia. It occupies an area of 2,297 km², of a total of approximately 184,920 km² of the Syrian Arab Republic. The region is bordered by the Mediterranean Sea to the west, the Anti-Lebanon Mountains to the south, Al Ghab Valley and the El-Asi (Orontes) River to the east and the Nur-Dağ Mountains to the north. It is composed of two subunits, each with distinctive physical-geographical, soil and climate features: continental Lattakia (Lattakia Mountains) (also named Jabal an Nusayryah or Alawite Mountains) and maritime Lattakia (HARDENBERG, 2003).

The continental subunit is represented by Jabal an Nusayryah Mountains. Between 0 and 60 m altitude, in the east, it has a geological structure dominated by sand, rocks sand, limestone and conglomerate. From 60 m to 200 m, the geological structure is of the Paleogene and Neogene age, composed of limestone, marl and clay. A filitic complex of rocks occurs in contact with the Alkabir Alshimali River (serpentine, peridotite, radiolarians, gabbro). The mountain slopes are covered mainly by forests of oak and cedar.

The maritime subunit belongs to the Levantine Basin of the Mediterranean Sea. It is located in the eastern Mediterranean, or the *Mare mediterraneum* (in Latin) that means the "sea in the middle of the land". This basin is the largest (2,969,000 km²) and deepest (average 1,460 m, maximum 5,267 m) enclosed sea on Earth (AL-AZKI, 2002; 2006; 2008).

MATERIAL AND METHODS

During the research carried out between 1985 and 2014, different stations were established for field observations, identification of species and collecting of material from the region.

For species identification, there were widely used field observations and photographs taken by us, during the field trips. Specimens were taken from marine sedimentary deposits belonging to the Cretaceous Period, of carbonate rocks (limestone and dolomite).

They were then examined in the laboratory of "Dr. Fawaz Azki" Geological Museum, Kismin, Lattakia - Syria. Fossil gastropods were determined by means of binocular magnifier and specimens of each genus were counted.

The nomenclature of taxa and data processing is according to the data from literature as well as to national and international legislation.

RESULTS AND DISCUSSIONS

Lattakia Region (33°36'55.02"N, 36°44'39.71"E) (Figs. 1; 2), part of the Middle East, is distinguished by spectacular landscape beauty and is characterized by rich and diverse assemblage, with many rare, endemic, relict species specific to the Levantine Basin and Western Arabian Province, and species on the edge of their geographical area, actual and fossil species. Its particular scientific significance is also derived from its geological, geomorphological, and paleontological characteristics (AXINI, 2011; 2012).



Figure 1. Earth map (from Google Earth, accessed: March 24, 2014) emphasizing the position of Lattakia Region in Asia.

Lattakia Region hosts many species of invertebrates and vertebrates fossils, of which, so far, we have identified over 135 species and genera, belonging to 4 classes and 2 phyla of invertebrates (Trilobita – Arthropoda; Gastropoda, Bivalvia, Cephalopoda – Mollusca) and 3 classes and 1 phylum of vertebrates (Osteichthyes, Reptilia, Mammalia – Chordata) (AL-AZKI 2012b, 2012c, 2012d; AL-AZKI & AXINI 2012a, 2012b; AL-AZKI, 2013; AL-AZKI & AXINI, 2013; AL-AZKI & AL-AZKI, 2014). Of fossil molluscs, gastropods occupy the first place, followed by ammonites and bivalves.

Of the fossil gastropods identified so far, the 16 genera include taxa among which 16 families grouped in 5 orders (AL-AZKI 2010-2011a, 2010-2011b; 2012a; BEGG & GRANT-MACKIE, 2003; BOUCHET & al., 2005; COX, 1960; COX & KNIGHT, 1960; DAYRAT & TILLIER, 2002; GARDNER, 1945; KNIGHT, 1956; MALAQUIAS & REID, 2008; MONTFORT, 1810; PONDER & LINDBERG, 1996, 1997; POPPE & TAGARO 2006; SCHMIDT & BELLEC, 1994; STEWART, 1927; TRACEY et al., 1993; WHITE, 1877) (Table 1).

The fossil gastropods genera were taken from sedimentary deposits belonging to the Cretaceous (AUBRY et al., 2009).



Figure 2. Geographical position of Lattakia Region (in the east of Cyprus, Nicosia) and with Palestine (in northern Jerusalem and in the northeast of Amman) (image from Google Earth).

Table 1. The taxonomic position of identified fossil gastropods.

Order	Family	Genus
Archaeogastropoda (Bellerophontida)?	Bellerophontidae	<i>Bellerophon</i> sp. (Montfort 1808)
Archaeogastropoda	Pleurotomariidae	<i>Pleurotomaria</i> sp. (Sowerby 1821)
Caenogastropoda (?Hypsogastropoda, ?Littorinimorpha)	Naticidae	<i>Natica</i> sp. (Scopoli 1777)
	Buccinidae	<i>Neptunea</i> sp. (Röding 1798) (syn. <i>Chrysodomus</i> sp. Swainson 1840, <i>Costaria</i> sp. Golikov 1977)
	Conidae	<i>Conus</i> sp. (Linnaeus 1758)
	Hipponicidae	<i>Hipponix</i> sp. (Defrance 1819)
	Cypraeidae	<i>Umbilia</i> sp. (Jousseau 1884)
	Viviparidae	<i>Viviparus</i> sp. (Montfort 1810)
	Cassidae	<i>Cassis</i> sp. (Scopoli 1777) (syn. <i>Cassidea</i> Bruguière 1789)
	Pachychilidae	<i>Brotia</i> sp. (H. Adams 1866) (syn. <i>Antimelania</i> P. Fischer & Crosse, 1892)
	Strombidae	<i>Rimella</i> sp. (Agassiz 1841)
	Ficidae	<i>Ficopsis</i> sp. (Conrad 1866)
	Pseudomelaniidae	<i>Bourguetia</i> sp. (Deshayes 1871)
Opisthobranchiata	Bullidae	<i>Bulla</i> sp. (Linnaeus 1758)
Euomphalina	Euomphalidae	<i>Euomphalus</i> sp. (J. De C. Sowerby 1814) (syn. <i>Phymatifer</i> , <i>Schizostoma</i> , <i>Paromphalus</i>)
Neritimorpha	Neritidae	<i>Velates</i> sp. (Montfort 1810)

In the Mesozoic Era, there occurred the breaking of the megacontinental landmass, Pangaea, into two supercontinents, Laurasia and Gondwana and the formation of the Tethys Ocean. From the Jurassic to the Miocene, this ocean separated the two supercontinents. Fragmentation of Gondwana by the Early Miocene allowed the annexation of India to Asia. This led to the closure of the Indo-Mediterranean-Atlanto Seaway, in parallel with the annexation of Africa to Europe. These tectonic events led to the isolation of the Mediterranean basin from the rest of the global ocean, and therefore isolation of its biodiversity (KWEAKASON, 2008). This has affected the species of gastropods, such as those of the genera *Bulla*, *Conus*, *Natica* (DEUSSEN, 1924; EDDIE'S, 2014; HALL, 2002; HOCUTT, 1987; NCBI, 2014; TAVIANI, 2002).

Umbilia sp. is a genus largely disappeared today. The 4 existing species present in Australian waters are cold water species (DARRAGH, 2002; WILSON & CLARKSON, 2004). This indicates that during the Cretaceous, there was a time when the Thetis Ocean presented ecological conditions similar to those of the current cold regions.

The current species of the Fam. Neritidae live in saltwater, freshwater, marine water, or brackish water as amphibious, either on rocks or mangrove trees. This demonstrates that the species of *Velates* genus lived in the Thetis Ocean or led an amphibious life on rocks or mangrove trees from stagnant or flowing waters on the shores of the ocean.

The species of the *Viviparus* genus lived in mud or on the macrophytes of stagnant or flowing terrestrial waters. The presence of fossil gastropods of this genus, identified by us in the Cretaceous deposits in Lattakia Region, indicates that the species of this genus, in that period, lived in mud or on macrophytes from stagnant waters on the Thetis Ocean shore or in its tributary rivers (GAN & DINELAY, 1997; GLAUBRECHT, 2010; NEGREA, 2002; DRABETT, 2012; STRAUS, 1996).

Today, the species of *Brotia* occur in freshwater habitats of Southeast Asia, ranging from Northern India in the west through to Sumatra in the east (India, Myanmar, Bangladesh, Thailand, Laos, Vietnam, China, Cambodia, Malaysia and Indonesia - Sumatra and Borneo only). They are found predominantly in fast flowing, well oxygenated rivers, also, occasionally in lakes. The presence of this genus of gastropods in Cretaceous deposits in Syria indicates that, at that time *Brotia* species were living in swift and well-oxygenated waters flowing into the Thetis Ocean (ADAMS, 1866; GAN & DINELAY, 1997; GLAUBRECHT, 2010; KÖHLER & DEIN, 2010; KÖHLER & GLAUBRECHT, 2001, 2006; KÖHLER et al., 2009, 2010; RABETT, 2012; STRAUS, 1996).

Gastropod populations occupy an important place in the aquatic biocoenosis structure, by the specific and numerical diversity and biomass (CHIRIAC & UDRESCU, 1965). They are among the first organisms to be investigated when examining waters. They indicate the oxygen content of the water of these basins. Their occurrence in surface water shows low levels of oxygen in water. The presence of these gastropods, in Cretaceous, shows us a high biodiversity in this geological period.

Also, the analysis of fossil gastropods species identified until now in this region indicates that the region climate in the Cretaceous was-like that of the current tropical areas and the Thetis Ocean waters, that covered the palaeo-biogeographical area in the Cretaceous period, presented ecological conditions like those of the current tropical

ocean waters. Predatory gastropods were benthic species, such as those of the genus *Conus* and *Natica*. They had a maximum development during this period. Prosobrachiates, such as those of the genera *Natica*, *Bellerophon*, *Euomphalus* were first and developed a number of marine types existing now (except *Bellerophon* and *Euomphalus*) (BRANSON, 1948; GILLULY, 1932; SMITH & CALKINS, 1904; THIRIOT-QUIEVREUX, 1983; WELLER, 1898; WESTGATE & KNOFF, 1932). All studied genera showed a high diversity in that period; that is indicated by the presence of multiple specimens in the sampling areas.

During the Cretaceous geological periods and until now, these gastropods contributed - by calcium carbonate present in the composition of their shells, to form the Cretaceous calcareous sedimentary rocks that make up largely this biogeographical region unique due to its natural history, geological structure, and palaeo-biodiversity.

While the Mediterranean Sea is a marine biodiversity hot spot, the Levantine Basin biodiversity is very little known (COLL et al., 2010).

It is well known that the current territory of Syria is the “cradle” of civilization. Here, after its appearance, the man ceased to migrate and started the first human settlements and for the first time here, he domesticated animals and cultivated wheat, invented the first alphabet and musical notes ... and the examples may continue. For this reason, this territory was strongly affected by human evolution from the beginning, with consequences on its biodiversity.

The actual communication between the Mediterranean, through the Levantine Basin, on the one hand, and the Red Sea and the Indo-Pacific Ocean, on the other hand - a communication created by man through the Suez Canal, created the possibility of many Indo-Pacific species migrating in the Levantine Basin. Its powerful damage due to invasive species has made the Levantine basin to be endangered. This can lead to the disruption of gastropods species existing in the Levantine Basin.

The analysis of the geographical elements of fossil gastropods from Lattakia Region (DEDIU, 2010a, 2010b, 2010c; PÂRVU, 2005) shows that the “Extinct in the wild” elements dominate with 7 taxa, followed by “Tethyan Relict” elements with 4 taxa and “Extinct in the region” elements with 3 taxa, and then “Tethyan Paleoendemit”, Southeast Asian, Indo-Pacific, Palaeartic, “not-evaluated in the region” elements, each with 1 taxon (Table 2).

Analysing the geological periods, we see that most genera appeared in the Mesozoic Era: Cretaceous – Pliocene (7 taxa), followed by Jurassic – Cretaceous age and Jurassic – Pliocene (by 2 taxa), and Cretaceous – Oligocene, Cretaceous – Miocene (by 1 taxon). 4 taxa appeared in the Palaeozoic Era: Early Cambrian - Early Cretaceous, Early Ordovician - Late Cretaceous, Carboniferous – Eocene, Carboniferous – Pliocene (NASSER, 2014).

Table 2. Comparative data on zoogeographical elements, ecological data and geological age of gastropods from Lattakia Region.

Genera	Zoogeoelements (original)	Actual ecological data/palaeo-ecological data (original)	Geological Age (original)
<i>Bellerophon</i> sp.	EW	P, SW, H, T, ST, OOS, PE, CA	ECA – EC
<i>Bourguetia</i> sp.	EW	EF	J – C
<i>Brotia</i> sp.	SA, ER	L, RW, SW, WO, VI	C – P
<i>Bulla</i> sp.	TR	SW, CL, T, PE, TI, C, N, HE, AL, PH, PS, TI	CAR – P
<i>Cassis</i> sp.	IP, ER	SW, T, TE, IT, H, PS, N, PR, STI	C – P
<i>Conus</i> sp.	TPE	SW, T, TI, IT, STI, WW, DW, PR, VE, CAR, EF, LI, PS, CA	C – P
<i>Euomphalus</i> sp.	EW	SW, STI, CA, DW, CL, CA, STI, PS, PER, EF, TER	EOR - LC
<i>Ficopsis</i> sp.	EW	SW, EF, STI, CAR, CL, TI, CA	C - M
<i>Hipponix</i> sp.	TR	SW, T, TE, CL, TI, EF, CA	C - P
<i>Natica</i> sp.	TR	SW, T, AMW, PR, EF	C - P
<i>Neptunea</i> sp.	TR	SW, T, IT, AMW, OM, PR, SC	J – P
<i>Pleurotomaria</i> sp.	EW	SW, T, DW, EF, PR	J – C
<i>Rimella</i> sp.	EW	SW	C - O
<i>Umbilia</i> sp.	ER	SW, T, STI, DW, CW, EF, PR, CAR	C - P
<i>Velates</i> sp.	EW	SW, L, B, EF, PH, LI, A, SAW, CL, TI, EU, FS	CAR - E
<i>Viviparus</i> sp.	P, NE	L, RW, PE, PH, EF	J – P

Abbreviations used:

Zoogeoelements: EW - Extinct in the wild; ER - Extinct in the region; SA - Southeast Asian; IP - Indo-Pacific; TR - Tethyan Relict (in the region); TPE - Tethyan Palaeoendemit (in region); P - Palaeartic; NE - not-evaluated in the region.

Actual ecological data/palaeo-ecological data (AXINI & AL-AZKI, 2012; WoRMS, 2014): A - amphibious; AL - algaophilous; AMW - arctic marine waters; CA - calciphilous; H - holoplanktonic snails; OOS - open oceans and seas; P - primitive snails; PE - pelagic living; ST - subtropical waters; SW - seawater snails; T - tropical waters; EF - epifaunal snails; L - freshwaters, VI - viviparous; RW - running waters; B - brackish, PE - pelophilous; PH - phytophilous, LI - lithophilous; SW - stagnant waters; WO - well oxygenated rivers; TI - tidal zones; IT - intertidal zones; STI - subtidal zones; PER - peritidal; N - nocturnal; HE - herbivorous; CL - coastlines; PS - psammophilous; TE - temperate waters; PR - predatory snails; VE - venomous; CAR - carnivorous; DW - bathyal zone; TER - terrestrial; SC - “scavenger” snails; OM - omnivores; CW - cool-waters; WW - warm-waters; SAW - salt waters; EU - eurybiont; FS - factor in speciation.

Geological Age: ECA - Early Cambrian; EOR - Early Ordovician; CAR - Carboniferous; J - Jurassic; EC - Early Cretaceous; C - Cretaceous; LC - Late Cretaceous; E - Eocene; O - Oligocene; M - Miocene; P - Pliocene.

CONCLUSIONS

The data presented in this work were compiled from field and laboratory studies conducted in 1985 - 2014.

Currently, there is no research on the palaeo-diversity of Latakia Region and Syria, in general. Our findings are unique. In the future, they will be upgraded with new data and research of laboratory and field. Also, the Levantine Basin biodiversity is very little known.

Moreover, our research on fossil species - plants and animals (invertebrates and vertebrates) are singular. No literature data were found in this field in Syria.

A total of 16 genera of fossil gastropods was identified in this time. Of these Mollusca, seven are extinct in the wild, three are extinct in the region, four are relict and one is a palaeo-endemit. Most genera appeared in the Mesozoic Era and they are marine snails.

The fossil and actual species that have been discovered in this area gives us the possibility of palaeo-environments reconstitution in this region (and also in the whole Syria) correlated with current environments, and also of the natural history of extinct and existing species of mollusks in Lattakia.

These fossils are part of the mollusks collections of the "Dr Fawaz Azki" Geological Museum, Kismin - Lattakia, Syria.

At present, there are many sites in this region which does not have a designated conservation status. Human impacts on different aspects may contribute to future declines and even disappearance of some gastropods (fossils or actual) in this bio-geographical region, many important species to science and human well-being, some of which have not been described yet.

REFERENCES

- ADAMS H. 1866. Description of a new genus and a new species of mollusks. *Proceedings of the Zoological Society of London*. London: 150-151.
- AL-AZKI F. 2002. *Geological Encyclopedia*. Hawaran Publisher Damascus. Syria. **1**. 1050 pp. (in Arabic).
- AL-AZKI F. 2006. *Geological Encyclopedia*. Hawaran Publisher Damascus. Syria. **2**. 1050 pp. (in Arabic).
- AL-AZKI F. 2008. *Geomorphology*. Syrian Arab Republic, Ministry of High Education, Tishreen University, Faculty of Science, Dep. of Geology. Tishreen. 300 pp. (in Arabic).
- AL-AZKI F. 2010-2011. *Geological Survey Field Geology*. Syrian Arab Republic, Ministry of High Education, Tishreen University, Faculty of Science, Dep. of Geology. 330 pp. (in Arabic).
- AL-AZKI F. 2010-2011. *Geological Survey Field Geology*. Syrian Arab Republic, Ministry of High Education, Tishreen University, Faculty of Science, Dep. of Geology. Tishreen. 240 pp. (in Arabic).
- AL-AZKI F. 2012. *Remote sensing*. Ministry of High Education, Tishreen University, Faculty of Science, Department of Geology. Tishreen. 210 pp. (in Arabic).
- AL-AZKI F. 2012. Correlation between marine fossils and biological data from Mediterranean Sea. In: Axini M. (ed.), 2012. *International Symposium Biodiversity Conservation "in situ" and "ex situ". First Edition. Volume. Program and abstracts*. Ex Ponto Publisher. Constanța: 38.
- AL-AZKI F. 2012. Determination of paleo-biodiversity and fossils. In: Axini M. (ed.), 2012. *International Symposium Biodiversity Conservation "in situ" and "ex situ". First Edition. Volume. Program and abstracts*. Ex Ponto Publisher. Constanța: 39.
- AL-AZKI F. 2012. The Dr. Fawaz Azki Geological Museum. In: Axini M. (ed.), 2012. *International Symposium Biodiversity Conservation "in situ" and "ex situ". First Edition. Volume. Program and abstracts*. Ex Ponto Publisher. Constanța: 69.
- AL-AZKI F. & AXINI MONICA. 2012a. Animal fossil collection of Dr. Fawaz Azki Geological Museum. In: Murariu D., Adam C., Chișamera G., Iorgu E., Popa O., Popa L. (eds). *Annual Zoological Congress of "Grigore Antipa" Museum – Book of abstracts*. Grigore Antipa National Museum of Natural History. Bucharest: 166.
- AL-AZKI F. & AXINI MONICA. 2012b. Animal fossil collection of Dr. Fawaz Azki Geological Museum. In: Murariu D., Adam C., Chișamera G., Iorgu E., Popa O., Popa L. (eds). *Annual Zoological Congress of "Grigore Antipa" Museum – Book of abstracts*. Grigore Antipa National Museum of Natural History. Bucharest: 166. Available online at: www.czga.ro/.../CZGA_2012_Book_of_Abstracts.pdf (Accessed: March 3, 2014).
- AL-AZKI F. 2013. Syria was home or passageway for dinosaurs at cretaceous period? In: Axini M., Al-Azki F. (eds.), 2013. *International Symposium Biodiversity Conservation "in situ" and "ex situ". Second Edition. Volume. Program and abstracts*. Ex Ponto Publisher. Constanța: 26.
- AL-AZKI F. & AXINI MONICA. 2013. Culture between geology and biology. In: Axini M., Al-Azki F. (eds.), 2013. *International Symposium Biodiversity Conservation "in situ" and "ex situ". Second Edition. Volume. Program and abstracts*. Ex Ponto Publisher. Constanța: 46.
- AL-AZKI (AXINI) MONICA & AL-AZKI F. 2014. "Dr. Fawaz Azki" Geological Museum, Kismin – Lattakia, Syria – World Heritage; Its Role In The Development Of Science And Environmental Education Worldwide. *Oltenia. Studii și comunicări. Științele Naturii*. Muzeul Olteniei Craiova (under press).

- AXINI MONICA. 2011. Research and Conservation of Biodiversity In: The Black Sea Romanian Coastal Zone. *Protection and Sustainable Management of the Black Sea Ecosystem, Third Millenium Imperative*. Book of Abstracts. Constanța: 75-76.
- AXINI MONICA. 2012. Gastropods Diversity In Conacu-Negrești Valley. *Oltenia. Studii și comunicări. Științele Naturii*. Muzeul Olteniei Craiova. **28**: 187-193.
- AXINI MONICA & AL-AZKI F. 2012. Biogeographical data on mollusks species in Dobrogea, Romania. In: Popescu O., Enache M., Zamfir M., Ștefănuț S. (eds.). *A New Approach of the Academic Research in Biology*. Romanian Academy. Institute of Biology Bucharest. Edit. Ars Docendi. Bucharest: 89-90.
- AUBRY M.-P., VAN COUVERING J. A., CHRISTIE-BLICK N., LANDING E., PRATT B. R., OWEN D. E., FERRUSQUÍA-VILLAFRANCA I. 2009. Terminology of geological time: Establishment of a community standard. *Stratigraphy*. Springer. London. **6**(2): 100-105.
- BEGG J. G. & GRANT-MACKIE J. A. 2003. New Zealand and New Caledonian Triassic Pleurotomariidae (Gastropoda, Mollusca). *Journal of the Royal Society of New Zealand*. Wellington. **33**: 223-268.
- BRANSON C. C. 1948. *Bibliographic index of Permian Invertebrates*. Geological Society of America Memoir. New York. **26**. 1049 pp.
- BOUCHET P. & ROCROI J.-P. (Ed.); Frýda J., Hausdorf B., Ponder W., Valdés Á. & Warén A. 2005. Classification and nomenclator of gastropod families. *Malacologia: International Journal of Malacology*. **47**(1-2). ConchBooks: Hackenheim. 397 pp.
- CHIRIAC E. & UDRESCU M. 1965. *Ghidul Naturalistului în Lumea Apelor Dulci*. Edit. Științifică. București: 126-141.
- COLL MARTA, PIRODDI CHIARA, STEENBEEK J., KASCHNER K., BEN RAIS LASRAM FRIDA, AGUZZI J., BALLESTEROS E., BIANCHI C. N., CORBERA J., DAILIANIS T., DANOVARO R., ESTRADA MARTA, FROGLIA C., GALIL B. S., GASO J. M., GERTWAGEN R., GIL J., GUILHAUMON F., KESNER-REYES KATHLEEN, KITSOS M.-S., KOUKOURAS A., LAMPADARIOU N., LAXAMANA E., DE LA CUADRA C. M. L. P. F., LOTZE H. K., MARTIN D., MOUILLOT D., ORO D., RAICEVICH S., RIUS-BARILE JOSEPHINE, SAIZ-SALINAS J. I., SAN VICENTE C., SOMOT S., TEMPLADO J., TURON X, VAFIDIS D., VILLANUEVA R., COLL ELENI V. M., PIRODDI CHIARA, JEROEN STEENBEEK, KRISTIN KASCHNER, FRIDA BEN RAIS LASRAM, JACOPO AGUZZI, ENRIC BALLESTEROS, CARLO NIKE BIANCHI, JORDI CORBERA, THANOS DAILIANIS, ROBERTO DANOVARO, MARTA ESTRADA, CARLO FROGLIA, BELLA S. GALI, JOSEP M. GASOL, RUTHY GERTWAGEN, JOAˆO GIL, FRANCˆOIS GUILHAUMON, KATHLEEN KESNER-REYES, MILTIADIS-SPYRIDON KITSOS, ATHANASIOS KOUKOURAS, NIKOLAOS LAMPADARIOU, ELIJAH LAXAMANA, CARLOS M. LO´PEZ-FE´ DE LA CUADRA, HEIKE K. LOTZE, DANIEL MARTIN, DAVID MOUILLOT, DANIEL ORO, SASˆA RAICEVICH, JOSEPHINE RIUS-BARILE, JOSE IGNACIO SAIZ-SALINAS, CARLES SAN VICENTE, SAMUEL SOMOT, JOSE´ TEMPLADO, XAVIER TURON, DIMITRIS VAFIDIS, ROGER VILLANUEVA, ELENI VOULTSIADOU MARTA COLL, CHIARA PIRODDI, JEROEN STEENBEEK, KRISTIN KASCHNER, FRIDA BEN RAIS LASRAM, JACOPO AGUZZI, ENRIC BALLESTEROS, CARLO NIKE BIANCHI, JORDI CORBERA, THANOS DAILIANIS, ROBERTO DANOVARO, MARTA ESTRADA, CARLO FROGLIA, BELLA S. GALI, JOSEP M. GASOL, RUTHY GERTWAGEN, JOAˆO GIL, FRANCˆOIS GUILHAUMON, KATHLEEN KESNER-REYES, MILTIADIS-SPYRIDON KITSOS, ATHANASIOS KOUKOURAS, NIKOLAOS LAMPADARIOU, ELIJAH LAXAMANA, CARLOS M. LO´PEZ-FE´ DE LA CUADRA, HEIKE K. LOTZE, DANIEL MARTIN, DAVID MOUILLOT, DANIEL ORO, SASˆA RAICEVICH, JOSEPHINE RIUS-BARILE, JOSE IGNACIO SAIZ-SALINAS, CARLES SAN VICENTE, SAMUEL SOMOT, JOSE´ TEMPLADO, XAVIER TURON, DIMITRIS VAFIDIS, ROGER VILLANUEVA, ELENI VOULTSIADOU. 2010. The Biodiversity of the Mediterranean Sea: Estimates, Patterns, and Threats. *Mediterranean Marine Diversity*. London. **5**(8): 1-36.
- COX L. R. 1960. Thoughts on the classification of the Gastropoda. *Proceedings of the Malacological Society of London*. London. **33**(6): 239–261.
- COX L. R. & KNIGHT J. B. 1960. Suborders of the Archaeogastropoda. *Proceedings of the Malacological Society of London*. London. **33**(6): 262–264.
- DAYRAT B. & TILLIER S. 2002. Evolutionary relationships of euthyneuran gastropods (Mollusca): a cladistic re-evaluation of morphological characters. *Zoological Journal of the Linnean Society*. London. **135**(4): 403-470.
- DARRAGH TH. 2002. A Revision of The Australian Genus *Umbilia* (Gastropoda: Cypraeidae). *Memoirs of the Museum of Victoria*. London. **59**(2): 355-392.
- DEDIU I. I. 2010. *Enciclopedie de ecologie*. Edit. Știința. Chișinău. 836 pp.
- DEDIU I. I. 2010. *Tezaurul terminologic al ecologiei*. Edit. Știința. Chișinău. 284 pp.
- DEDIU I. I. 2010. *Axiomatica. Principiile și legile ecologiei*. Edit. Știința. Chișinău. 216 pp.
- DEUSSEN A. 1924. *Geology of the Coastal Plain of Texas West of Brazos River*. U. S. Government Printing Office. London. **126**. 139 pp.
- GAN J. Y & DINELAY D. L. 1997. Palaeontology and Historical Geology. *Proceedings of the 30th International Geological Congress*. Beijing. **12**: 1-192.

- GARDNER JULIA 1945. *Mollusca of the Tertiary Formations of Northeastern Mexico*. Geological Society of America. Memoir 11. New York. 331 pp.
- GILLULY J. 1932. Geology and Ore Deposits of the Stockton and Fairfield Quadrangles, Utah. *United States Geological Survey, Geological Survey professional paper*, U. S. Government Printing Office. Utah. **173**: 12-230.
- GLAUBRECHT M. 2010. *Evolution in Action: Case studies in Adaptive Radiation, Speciation and the Origin of Biodiversity*. Biomedical and Life Science. Springer. 611 pp.
- HALL C. A. 2002. *Nearshore Marine and Paleoclimatic Regions, Increasing Zoogeographic Provinciality, Molluscan Extinctions, and Paleoshorelines, California. Late Oligocene (27 Ma) to Late Pliocene (25 Ma)*, edition 357, Geological Society of America. New York. **357**. 489 pp.
- HARDENBERG M. F. 2003. *Tectonics and sedimentation of early continental collision in the Eastern Mediterranean (Northwest Syria)*. Thesis submitted for the degree of Doctor of Philosophy. University of Edinburgh. 366 pp.
- HOCUTT CH. H. 1987. Evolution of the Indian Ocean and the drift of India: A vicariant event. *Hydrobiologia*. Elsevier. Stuttgart. **150**(3): 203-223.
- KNIGHT J. B. 1956. New families of Gastropoda. *Journal of the Washington Academy of Sciences*. Washington. **46**(2): 41-42.
- KÖHLER F. & DEEIN G. 2010. Hybridisation as potential source of incongruence in the morphological and mitochondrial diversity of a Thai freshwater gastropod (Pachychilidae, *Brotia* H. Adams, 1866). *Zoosystematics and Evolution*. Wiley-VCH Verlag. Stuttgart. **86**(2): 301-314.
- KÖHLER F., DU L.-N., YANG J.-X. 2010. A new species of *Brotia* from Yunnan, China (Caenogastropoda, Pachychilidae). *Zoosystematics and Evolution*. Wiley-VCH Verlag. Stuttgart. **86**(2): 295-300.
- KÖHLER F. & GLAUBRECHT M. 2001. Toward a systematic revision of the Southeast Asian freshwater gastropod *Brotia* H. – Adams, 1866 (Cerithioidea: Pachychilidae): an account of species from around the South China Sea. *Journal of Molluscan Studies*. Wiley-VCH Verlag. Stuttgart. **67**(3): 281-318.
- KÖHLER F. & GLAUBRECHT M. 2006. A systematic revision of the Southeast Asian freshwater gastropod *Brotia* (Cerithioidea: Pachychilidae). *Malacologia*. Elsevier. Stuttgart. **48**: 159-251.
- KÖHLER F., HOLFORD DO V. T., HO T. H. 2009. Exploring a largely unknown fauna: On the diversity of pachychilid freshwater gastropods in Vietnam (Caenogastropoda: Cerithioidea). *Molluscan Research*. Magnolia Press. London. **29**(3): 121-146.
- KWEAKASON P. M. 2008. *A Holocene Prehistoric Sequence in the Egyptian Red Sea Area: The Tree Shelter*. In: Egyptian Prehistory monographs. Leuven University Press. Leuven. **7**. 104 pp.
- MALAQUIAS M. A. & REID D. G. 2008. Systematic revision of the living species of Bullidae (Mollusca: Gastropoda: Cephalaspidea), with a molecular phylogenetic analysis. *Zoological Journal of the Linnean Society*. London. **153**: 453-543.
- MONTFORT D. DE. 1810. *Conchyliologie systématique, et classification méthodique des coquilles; offrant leurs figures, leur arrangement générique, leurs descriptions caractéristiques, leurs noms; ainsi que leur synonymie en plusieurs langues. Ouvrage destiné à faciliter l'étude des coquilles, ainsi que leur disposition dans les cabinets d'histoire naturelle*. Coquilles univalves, non cloisonnées. Paris. **2**. 676 pp.
- NASSER H. 2014. Fossils - "Words" In "Sedimentary Rocks Book" Which "Describe Us" The Life History On The Earth. *RJGB*. Washington. **3** (under press).
- NEGREA ALEXANDRINA. 2002. Class Gastropoda. Melci. In: Godeanu (Ed.). Diversitatea lumii vii. *Determinatorul Ilustrat al Florei și Faunei României*. Vasile Goldiș University Press. Arad. **2**: 338-343.
- PÂRVU C-TIN. (coord.). 2005. *Dicționar enciclopedic de mediu*. Regia Autonomă MONITORUL OFICIAL. București. **1**(A-K). 852 pp., **2**(L-Z). 777 pp.
- PONDER W. F. & LINDBERG D. R. 1996. Gastropod phylogeny—challenges for the 90s. In: Taylor J. (ed.) *Origin and Evolutionary Radiation of the Mollusca*. Oxford University Press. Oxford: 135–154.
- PONDER W. & LINDBERG D. R. 1997. Towards a phylogeny of gastropod molluscs: an analysis using morphological characters. *Zoological Journal of the Linnean Society*. Paris. **119**(2): 83-265.
- POPPE G. T. & TAGARO S. P. 2006. *The New Classification of Gastropods According to Bouchet & Rocroi, 2005*. Visaya. 11 pp.
- RABETT R. J. 2012. *Human Adaptation in the Asian Palaeolithic: Hominin Dispersal and Behaviour during the Late Quaternary*. Cambridge University Press. 372 pp.
- SCHMIDT W. & BELLEC O. 1994. Findings of some uncommon sea-shells off Madagascar. *African Journal of Tropical Hydrobiology and Fisheries*. **5**(1): 63 – 66.
- SMITH G. O. & CALKINS F. C. 1904. A Geological Reconnaissance Across the Cascade Range Near the Forty-ninth Parallel. *U. S. Geological Survey bulletin* (edition 235). U. S. Government Printing Office. New York. 103 pp.
- STEWART R. B. 1927. Gabb's California fossil type gastropods. *Proceedings of the Academy of Natural Sciences of Philadelphia*. **78** (for 1926): 287–447, pls. 20–32.
- STRAUS L. G. 1996. *Humans at the End of the Ice Age: The Archaeology of the Pleistocene – Holocene Transition*. Interdisciplinary Contributions to Archaeology Language of science. Springer. London. 376 pp.
- TAVIANI M. 2002. The Mediterranean Benthos From Late Miocene Up To Present: Ten Million Years Of Dramatic Climatic And Geologic Vicissitudes. *Biology Marine of Mediterranean*. Elsevier. London. **9**(1): 445-463.

- THIRIOT-QUIEVREUX CATHERINE. 1983. Summer Meroplanktonic Prosobranch Larvae Occuring off Beaufort, North Carolina. *Estuaries*. London. **6**(4): 387-398.
- TRACEY S., TODD J. A., ERWIN D. H. 1993. Mollusca, Gastropoda. In: M. J. Benton (ed.) *The Fossil Record*. Chapman & hall. London. **2**: 131–167.
- WELLER S. 1898. *A Bibliographic Index of North American Carboniferous Invertebrates*. U.S. Government Printing Office. New York. **153**(8). 653 pp.
- WESTGATE L. G. & KNOPF A. 1932. *Geology and Ore Deposits of the Pioche District, Nevada*. Geological Survey professional paper. U. S. Government Printing Office. **171**. 79 pp.
- WHITE C. A. 1877. *Report upon the invertebrate fossils collected in portions of Nevada, Utah, Colorado, New Mexico, and Arizona, by parties of the expeditions of 1871, 1872, 1873, and 1874*. Report upon United States geographical surveys west of the one hundredth meridian. Paleontology. Government printing Office. Washington. **4**. 219 pp., 21 plates.
- WILSON B. & CLARKSON P. 2004. *Australia's spectacular cowries. A review and field study of two endemic genera: Zoila and Umbilia*. Odyssey Publishing. El Cajon. California. 396 pp.
- ***. Eddie's Shell Catalogue. 2014. http://parallel.park.org/Guests/Shells/Shell_Catalogue/Shell_Pages/Shell_Catalogue_Home_Page.html (Accessed: March 6, 2014).
- ***. Fossilworks. Gateway to the Paleobiology Database. www.fossilworks.org (Accessed: March 6, 2014).
- ***. NCBI. 2014. Taxonomy Browser. [http://www.ncbi.nlm.nih.gov/Taxonomy/taxonomyhome.html/...](http://www.ncbi.nlm.nih.gov/Taxonomy/taxonomyhome.html/) (Accessed: March 6, 2014).
- ***. WoRMS. World Register of Marine Species. 2014. [http://www.marinespecies.org/...](http://www.marinespecies.org/) (Accessed: March 7, 2014).

Axini Monica, Al-Azki Fawaz

"Monachus" Group of Scientific Research and Ecological Education, 8, Hortensiei Alley, Constanța 900518, Romania.

"Dr Fawaz Azki" Geological Museum, Ain Al-Bayda, Kismis-Lattakia, Syria.

E-mail: monica.gceem@gmail.com, fawazazki@yahoo.com

Received: March 31, 2014
Accepted: June 7, 2014