ACADEMIA ROMÂNĂ INSTITUTUL DE ARHEOLOGIE "VASILE PÂRVAN"

MATERIALE ȘI CERCETĂRI ARHEOLOGICE

SERIE NOUĂ V

EDITURA ACADEMIEI ROMÂNE BUCUREȘTI, 2009

https://biblioteca-digitala.ro

Radiocarbon Dating and Faunal Stable Isotopes for the Galeria Principală, Peștera Muierii, Baia de Fier, Gorj County, Romania

ADRIAN DOBOŞ, ANDREI SOFICARU, AURELIAN POPESCU and ERIK TRINKAUS*

Key words: Upper Paleolithic, Middle Paleolithic, cave bears, Late Pleistocene, diet.

Abstract: Radiocarbon dates and stable isotope signatures are presented for six faunal samples from the Galeria Principală, Peştera Muierii. Despite stratigraphic incertainties, the samples should date from the earlier Upper Paleolithic of this portion of the site to ~30,000 ¹⁴C BP and the Middle Paleolithic levels to ~41,000 to ~47,500 ¹⁴C BP, with the deeper levels being beyond radiocarbon age (>52,000 ¹⁴C BP). These dates therefore imply that the Pleistocene human occupation of the site consists of a relatively late Early Upper Paleolithic (or very early Middle Upper Paleolithic) and a fairly late Middle Paleolithic. Stable isotope analysis of the dated samples reinforces the ecological flexibility of Late Pleistocene cave bears.

Cuvinte-cheie: Paleolitic superior, Paleolitic mijlociu, urși de peșteră, Pleistocen târziu, dietă.

Rezumat: În acest articol sunt prezentate datele radiocarbon și analizele izotopice obținute pentru șase eșantioane faunistice prelevate din Galeria Principală din Peștera Muierii. În ciuda provenienței stratigrafice incerte, ele datează nivelul paleolitic superior vechi din această zonă a peșterii la cca. 30,000 ¹⁴C BP, iar nivelurile caracteristice Paleoliticului mijlociu le situează între cca. 41,000 și 47,500 ¹⁴C BP. Aceste date arată că pentru Pleistocen, ocupația umană se plasează la sfîrșitul a paleoliticului superior vechi (sau chiar la începutul Paleoliticului superior mijlociu); în ceea ce privește paleoliticul mijlociu de aici, acesta este relativ târziu. Analizele izotopice pun în evidență flexibilitatea ecologică a urșilor de peșteră în Pleistocenul Superior.

Introduction

The archeological excavations in the Peştera Muierii (also, Peştera Muierilor), Baia de Fier, Gorj County (45° 11' N, 23° 46' E), initially in the

1920s and then more systematically in the early 1950s, yielded an abundance of Paleolithic and more recent archeological remains and a large sample of Pleistocene faunal remains (Nicolăescu-Plopsor 1935, 1956, 1957; Daicoviciu et al. 1953; Gheorghiu et al. 1954; Gheorghiu and Haas 1954; Nicolăescu-Plopșor et al. 1957). Even though the Paleolithic archeological remains continue to pertain to discussions of the Paleolithic of Romania (especially of the Middle Paleolithic and the early Upper Paleolithic) (Cârciumaru 1999; Păunescu 1989, 2000; Dobrescu 2008), it is principally the early modern human fossil remains from the site which have drawn attention (Gheorghiu and Haas 1954; Nicolăescu-Plopșor 1968; Soficaru et al. 2006; Cosac, 2006-07; Trinkaus, 2008). These human remains, directly radiocarbon dated to $\sim 30,000^{-14}$ C BP (~35,000 cal BP) (Olariu et al., 2003; Soficaru et al., 2006), have consequently joined the small sample of securely dated European human fossils that derive from the time period of the Early Upper Paleolithic (Trinkaus, 2005, 2007). The remainder of the contents of the complex cave system, however, remains poorly dated.

It is readily apparent from the existing fieldnotes and the collections in various institutions [to our knowledge, there are paleontological and/or archeological remains from the Pestera Muierii in at least the Muzeul Olteniei (Craiova), Institutul de Arheologie "Vasile Pârvan" (București), Institutul de Antropologie "Fr. J. Rainer" (București), Institutul de Speologie "Emil Racoviță" (București), Muzeul Național de Istorie (București), and Muzeul Militar National (București)] that any attempt to securely date the deposits of the Peştera Muierii would require the re-excavation of portions of the cave and the acquisition of datable material from primary contexts within stratigraphic profiles. This situation is especially evident in the existing collections,

MATERIALE ȘI CERCETĂRI ARHEOLOGICE (serie nouă) V, 2009, p. 15-20

^{*}Adrian Doboş: Institutul de Arheologie "Vasile Pârvan," Henri Coandă 11, sector 1, 010667 București, Romania; Andrei Soficaru: Institutul de Antropologie "Fr. J. Rainer," Eroii Sanitari 8, P.O. Box 35-13, București 050471, Romania; Aurelian Popescu: Muzeul Olteniei, Madona Dudu nr. 44, 200410 Craiova, Romania; Erik Trinkaus: Department of Anthropology, Campus Box 1114, Washington University, Saint Louis MO 63130, USA.

since they can only represent a partial retention of paleontological and archeological remains from the excavations, and none of the remains has detailed stratigraphic and horizontal proveniences other than indications of the gallery, trench, and depth below datum marked on a minority of the remains. Nonetheless, as part of the project to securely date and analyze the Muierii human remains (Soficaru *et al.*, 2006), samples of faunal remains from a stratigraphic sequence were dated to help provide a general chronological framework for the site's deposits. It is hoped that these dates, presented here, will assist in establishing a temporal framework for the Paleolithic deposits of the Peştera Muierii.

The Peştera Muierii

The Pestera Muierii consists of a series of interconnected galleries at multiple levels (cf., plans in Gheorghiu and Haas 1954; Păunescu 2000). The largest gallery, the Galeria Principală, is >70 m long and 5-10 m wide, with a smaller section near the entrance, the Gura Peşterii, being the south cave entrance. Parallel to it is a narrower gallery at a lower elevation, Galeria Secundară towards the front of the system and the Galeria Musteriană deeper within. This lower chamber connects with the Galeria Principală at the front of the Galeria Secundară and at the back of the Galeria Musteriană. Additional galleries extend off of these main passageways and at their ends away from the modern entrance. The principal excavations, carried out from 1951 to 1953, consisted of a trench across the Gura Peşterii, two trenches at right angles to each other in the entrance portion of the Galeria Principală (Sector I and Sector II, with Sector I being deeper within the cave), three small soundings deep within the Galeria Principală, and then most of the surface areas of the Galeria Secundară and the Galeria Musteriană (Gheorghiu and Haas 1954, fig. 13). The faunal dating samples all derive from Sector I of the Galeria Principală, excavated in 1952.

Previous Radiocarbon Dates for the Peştera Muierii

In addition to the series of radiocarbon dates on the human remains from the Peştera Muierii (Soficaru *et al.* 2006), there have been only two previous attempts to date the deposits of the cave system. A bone from the Mousterian Level I of Sector C of the Galeria Musteriană, at a depth of 1.40 - 1.50 m, provided an age of 42,560 +1,310, -1,120 ¹⁴C BP (GrN-16977) (Păunescu 2000). In addition, a bear bone (Ursus spelaeus?) of unknown provenience within the cave system and of unknown stratigraphic depth provided an age of >29,000 ¹⁴C BP (Mo-105) (Vinogradov et al. 1968). The first date may be either finite or a minimum age; if the former, it is a reasonable age for a relatively late phase of the Middle Paleolithic. The second date only serves to establish that there were the remains of bears at least of oxygen isotope stage 3 age within the cave system, a not surprising result.

New Faunal Dates for the Galeria Principală

Given the general absence of horizontal provenience for the faunal remains from the Peştera Muierii, we selected six samples, all labeled "B.F.52" (Baia de Fier 1952), "G.P." (Galeria Principală), "S.I" (Sector I), and then a depth or a depth range below datum (Table 1). Four of them are diaphyseal samples from metapodials of *U. spelaeus*, one is a diaphyseal sample from a metapodial of a large felid (*Panthera spelaea?*), and the last is a root of a maxillary molar of *Alces alces*.

The stratigraphic depths of the samples, if datum is near the surface of the original deposits and the schematic stratigraphic profile of Păunescu (2000: fig. 118) can be used as a general guide, should span the levels designated "Aurignacian" and "Mousterian" with the deepest sample possibly predating the Middle Paleolithic accumulation in this portion of the cave. This assumes that 1) the levels across Sector I of the Galeria Principală were reasonably horizontal (which is clearly not the case, as drawn), 2) the samples all derive from one part of Sector I, 3) Păunescu's stratigraphic profile is accurately to scale, and 4) the reference datum was close to the surface. At present, none of these assumptions are verifiable, but all of them should provide a working framework for the samples.

The samples were removed with a stainless steel disk saw on a Dremel tool, and submitted to the Oxford Radiocarbon Accelerator Unit (ORAU) for accelerator mass spectrometry (AMS) dating. None of the samples had macroscopic evidence of any form of preservative. All of the samples, being of bone and tooth root, were pretreated following standard ORAU procedure (Brown et al., 1988) combined with ultrafiltration (Higham et al. 2006). Ultrafiltration sorts the collagen fraction by molecular weight, removes smaller fractions likely to be from contamination (<30 kD), and dates only the higher (>30 kD) molecular weight portion (Higham *et al.* 2006). Ultrafiltration commonly

produces dates that are modestly older and/or more precise than AMS dates run without ultrafiltration.

Sample bone and root weights range from 520 mg to 900 mg, with collagen yields beween 12.0 and 77.5 mg (Table 1). Most importantly, the carbon-nigrogen (C:N) ratios are all between 2.9 and 3.5, as they should be for well preserved bone (DeNiro 1985). The δ^{13} C values for the *U. spelaeus* samples are moderately negative, but this is common for samples of *U. spelaeus* (Richards *et al.*, 2007, and references therein).

Table 1

Radiocarbon results for faunal remains from the Peştera Muierii, Baia de Fier, Romania. All samples are from the depth below datum brackets indicated from the Galeria Principală (GP), Sector I (SI), 1952 excavations. All samples were run using ultrafiltration (Higham 2006) at the Oxford Radiocarbon Accelerator Unit (ORAU). Calendrical years based on CalPal 2007 (www.calpal.de)

Sample number	Mui05-5	Mui05-6	Mui05-8	Mui05-9	Mui05-10	Mui05-11
Curating institution	Inst. Archeol.	Muzeul Olteniei	Muzeul Olteniei	Inst. Archeol.	Inst. Archeol.	Inst. Archeol.
GP SI depth (m below datum)	0.90	1.20 - 1.40	1.40 - 1.60	1.60 - 1.70	1.70 - 1.90	1.90 - 2.05
ORAU number	OxA-15554	OxA-15530	OxA-16380	OxA-16381	OxA-16382	OxA-16383
Species	Alces alces	Ursus spelaeus	Panthera spelaea	Ursus spelaeus	Ursus spelaeus	Ursus spelaeus
Bone	Molar root	Metapodial	Metapodial	Metapodial	Metapodial	Metapodial
Radiocarbon age (¹⁴ C years BP)	30,060 ± 280	$40,850 \pm 450$	$47,500 \pm 900$	40,950 ± 450	42,700 ± 550	>52,400
"Calendrical" age (cal years BP)	35,367 ± 318	$44,372 \pm 790$	$51,292 \pm 2081$	44,467 ± 783	46,182 ± 1176	
$\delta^{13}C$	-19.9‰	-20.3‰	-19.1‰	-20.3‰	-20.2‰	-20.7‰
C:N	3.5	3.3	3.2	3.2	3.2	3.2
Sample weight (mg)	520	640	900	800	840	800
Collagen weight (mg) (ultrafiltered gelatin yield)	13.1	52.2	77.5	51.7	60.5	12.0
Burnweight (mg) (gelatin combusted for graphitization)	5.6	5.3	5.4	5.3	5.2	5.2
%C (% carbon on combustion)	46.9%	43.4%	48.8%	47.5%	46.9%	44.9%
%N (% nitrogen on combustion)	16.1%	15.1%	17.9%	17.5%	17.2%	16.4%
$\delta^{15}N$	(7.3)	6.0	8.2	3.7	3.4	7.3

The resultant samples provide a series of dates that are bracketted by the A. alces molar at -0.90 m and ~ 30 ka ¹⁴C BP and a U. spelaeus metapodial at -1.90-2.05 m at >52.4 ka ¹⁴C BP. Three of the remaining four samples, between -1.20 and -1.90 m, cluster around 41 to 43 ka ¹⁴C BP, with one sample, the felid metapodial, being substantially older but still finite at ~47.5 ka ¹⁴C BP (Table 1). If the reference to the schematic chart of Păunescu (2000) has any validity, the most recent date is probably associated with the small early Upper Paleolithic assemblage, one which has been designated "Aurignacian" but typologically and technologically could as well derive from an early Gravettian level. The deepest and oldest date may well predate the Middle Paleolithic occupation of at least this portion of the cave. The other samples, despite either vertical mixing of the remains (always possible with smaller bones such as metapodials) or horizontal variance in the depths of the levels (likely given the horizontal extent of Sector I of the Galeria Principală), appear to date a relatively late phase of the Middle Paleolithic, well within OIS 3.

Carbon and Nitrogen Stable Isotopes

In addition to the radiocarbon dates, the ORAU sample preparation and analysis provide carbon $(\delta^{13}C)$ and nitrogen $(\delta^{15}N)$ stable isotopes for these samples (Table 1). The bone collagen isotope values likely reflect dietary protein consumed over the past 10 to 20 years of life for an adult longlived mammal (Wild et al., 2000). Although the δ^{13} C values can be used to assess whether the dietary protein carbon is derived from marine versus terrestrial sources or from C₃ or C₄ plants, it is the trophic level of the protein consumed that is of interest here. The $\delta^{15}N$ of bone collagen is ~2‰ to ~4‰ higher than dietary protein (Schoeninger and DeNiro 1984; Schwarcz and Schoeninger 1991), and one therefore can use it to see if dietary protein derived mainly from plants or from higher trophic level protein sources, especially as compared to other fauna from the same site. Of particular relevance here, since most of the samples are from U. spelaeus, is experimental work with living bears (Hilderbrand et al. 1996) that shows bone collagen δ^{15} N values to be an accurate reflection of diet. In general δ^{15} N values for herbivores are <6.0‰, >8.0‰ for carnivores, and in between for omnivores (Bocherens 2002).

Unfortunately only one specimen of a herbivore is available in this sample, the molar root of *A. alces*. Tooth roots (having been formed early in development) sometimes provide augmented δ^{15} N values relative to adult values (Jenkins *et al.* 2001). The value of 7.3 ‰ is therefore in parentheses, indicating that it may be anomalous (it is certainly high compared to most herbivore values) and should not be taken to represent the trophic level of this species.

The δ^{15} N value of 8.2 % for the large felid is within the carnivore range but relatively low for as pure a carnivore as felids tend to be. In contrast, the range of $\delta^{15}N$ values for the cave bears (3.4 to 7.3 ‰) spans much of the range documented for these large Late Pleistocene ursids (Richards et al. 2007). Isotope values have long been used to argue that cave bears were exclusively vegetarian (e.g., Bocherens, 2002), but recent data from the Peştera cu Oase (Caraş-Severin County) supplemented with data from Pestera Cioclovina and Pestera Muierii have shown that cave bears in fact span a range of dietary profiles, from purely vegetarian to moderately carnivorous (Richards et al. 2007), undoubtedly reflecting the same kinds of ecological and dietary flexibility known for extant brown (U. arctos) and black (U. americanus) bears (MacHutchon and Wellwood 2003; Belant et al. 2006; Mowat and Heard 2006). The four samples of U. spelaeus from the Peştera Muierii reinforce this pattern.

Conclusions

We remain a long way from sorting out the chronology and associated archeology and paleobiology of the Late Pleistocene occupants (human and faunal) of the Peştera Muierii. However, in addition to the early modern human remains from the site bringing the locality into paleoanthropological awareness, it may be possible through analyses of the preserved remains from the earlier excavations to shed some light on the site and its occupations by humans, bears, and other denizens. These radiocarbon determinations, despite uncertainties regarding their stratigraphic associations, provide additional data on the time span of the Pleistocene paleontological and anthropological deposits within the cave system. It is hoped that they will provide a further foundation for future interpretations of the site and its contents.

Acknowledgments

The sampling of the Peştera Muierii faunal remains in the Muzeul Olteniei and in the Institutul de Arheologie "Vasile Pârvan" was done with the permission of Drs. M.-F. Fifor and R. Dobrescu respectively. The analysis and dating of the Peştera Muierii remains has been funded by the National Science Foundation (BCS-0509072) and the Wenner-Gren Foundation (7290). To all of them we are grateful.

REFERENCES

- Belant et al. 2006 J. L. Belant, K. Kielland, E. H. Follmann, L. G. Adams, *Interspecific resource partitioning in sympatric* ursids, Ecological Applications 16, 2006, p. 2333–2343.
- Bocherens 2002 H. Bocherens, Alimentation des ours et signatures isotopiques, ERAUL 100, 2002, p. 41-49.
- Browns et al. 1988 T. A. Brown, D. E. Nelson, J. S. Vogel, J. R. Southon, *Improved collagen extraction by modified Longin method*, Radiocarbon 30, 1988, p. 171-177.
- Cârciumaru 1999 M. Cârciumaru, Le Paléolithique en Roumanie, Grenoble, 1999.
- Cosac 2006-2007 M. Cosac, Une réévaluation du contexte des découvertes d'ossements humains de la galerie M de la grotte Muierilor à Baia de Fier, Annales d'Université "Valahia" Târgovişte. Section d'Archéologie et d'Histoire 8-9, 2006-2007, p. 199-205.
- Daicoviciu et al. 1953 C. Daicoviciu, Ş. Ferenczi, A. Bodor, C.S. Nicolăescu-Plopşor, N. Gostar, D. Radu, M. Detiu, P. Duka, Date preliminare asupra rezultatelor paleoantropologice de la "Peştera Muierilor", R. Novaci, Reg. Craiova, SCIVA 1-2, 1953, p. 195-207.
- DeNiro 1985 M. J. DeNiro, Post-mortem preservation and alteration of in vivo bone collagen isotope ratios in relation to paleodietary reconstruction, Nature 317, 1985, p. 806-809.
- Dobrescu 2008 R. Dobrescu, *Aurignacianul din Transilvania*, București, 2008.
- Gheorghiu and Haas 1954 A. Gheorghiu, N. Haas, Date privind omul primitiv de la Baia de Fier. Considerații paleoantropologice, Sesiunea Secțiunii de Științe Medicale a Academiei R.P.R. 41, 1954, p. 641-63.
- Gheorghiu et al. 1954 A. Gheorghiu, C.S. Nicolăescu-Plopşor, N. Haas, E. Comşa, C. Preda, G. Bombiţă, G. Enea, F. Gheorghiu, S. Iofcea, C. S. Nicolăescu-Plopşor, D. Neagoe, R. Silveanu, Raport preliminar asupra cercetărilor de

paleontologie umană de la Baia de Fier (Reg. Craiova) din 1951, Probleme de Antropologie 1, 1954, p. 73-86.

- Haas 1957 H. Haas, Critica criteriului "Schwalbe" în determinarea scapulei de Homo primigenius, Probleme de Antropologie 3, 1957, p. 21-46.
- Higham et al. 2006 T. F. G. Higham, R. M. Jacobi, C. Bronk Ramsey, AMS radiocarbon dating of ancient bone using ultrafiltration, Radiocarbon 48, 2006, p. 179–195.
- Hilderbrand et al. 1996 G. V. Hilderbrand, S. D. Farley, C.T. Robbins, T. A. Hanley, K. Titus, C. Servheen, Use of stable isotopes to determine diets of living and extinct bears, Canadian Journal of Zoology 74, 1996, p. 2080-2088.
- Jenkins et al. 2001 S. G. Jenkins, S. T. Partridge, T. R. Stephenson, S. D. Farley, C. T. Robbins, Nitrogen and carbon isotope fractionation between mothers, neonates, and nursing offspring, Oecologia 129, 2001, p. 336–341.
- MacHutchon, Wellwood 2003 A.G. MacHutchon, D.W. Wellwood, *Grizzly bear food habits in the northern Yukon*, *Canada*, Ursus 14, 2003, p. 225-235.
- Mowat, Heard 2006 G. Mowat, D.C. Heard, *Major components* of grizzly bear diet across North America, Canadian Journal of Zoology 84, 2006, p. 473-489.
- Nicolăescu-Plopșor 1935 C. S. Nicolăescu-Plopșor, *Le Paléolithique en Roumanie*, Dacia 5, 1935, p. 41-107.
- Nicolăescu-Plopșor 1956 C. S. Nicolăescu-Plopșor, *Rezultatele principale ale cercetărilor paleolitice în ultimii patru ani în R.P.R.*, SCIV 1-2, 1956, p. 7-39.
- Nicolăescu-Plopșor 1957 C. S. Nicolăescu-Plopșor, Le Paléolithique dans la République Populaire Roumaine à la lumière des derniers recherches, Dacia 1, 1957, p. 41-60.
- Nicolăescu-Plopşor et al. 1957 C.S. Nicolăescu-Plopşor, E. Comşa, D. Bolomey, Şantierul arheologic Baia de Fier, Materiale 3, 1957, p. 13-26.
- Nicolăescu-Plopşor 1968 D. Nicolăescu-Plopşor, Les hommes fossiles découverts en Roumanie, in The VIIth International Congress of Anthropological and Ethnological Sciences, Moscow, 1968, p. 381-386.
- Olariu et al. 2003 A. Olariu, E. Alexandrescu, G. Skog, R. Hellborg, K. Stenström, M. Faarinen, P. Persson, Dating of two paleolithic human fossils from Romania by accelerator mass spectrometry in M. Oancea, C. Schiaua, D. Grecu, M. Dumitriu (eds.) IFIN-HH Scientific Report 2001-2002,: "Horia Hulubei" National Institute for Physics and Nuclear Engineering. Bucureşti, 2003, p. 81.
- Păunescu 1989 A. Păunescu, Le Paléolithique et le Mésolithique de Roumanie (un bref aperçu), L'Anthropologie 93, 1989, p. 123-158
- Păunescu 2000 A. Păunescu, Paleoliticul şi Mezoliticul din spațiul cuprins între Carpați şi Dunăre, Bucureşti, 2000.
- Richards et al. 2008 M. P. Richards, M. Pacher, M. Stiller, J. Quilès, M. Hofreiter, S. Constantin, J. Zilhão, E. Trinkaus, Isotopic evidence for omnivory among European cave bears: Late Pleistocene Ursus spelaeus from the Peştera cu Oase, Romania, Proceedings of the National Academy of Sciences USA 105, 2008, p. 600-604.
- Schoeninger, DeNiro 1984 M. Schoeninger, M. DeNiro, Nitrogen and carbon isotopic composition of bone

collagen from marine and terrestrial animals, Geochimica Cosmochimica Acta 48, 1984, p. 625-639.

- Schwarcz, Schoeninger 1991 H. Schwarcz, M. Schoeninger, Stable isotope analyses in human nutritional ecology, Yearbook of Physical Anthropology 34, 1991, p. 283-321.
- Soficaru et al. 2006 A. Soficaru, A. Doboş, E. Trinkaus, Early modern humans from the Peştera Muierii, Baia de Fier, Romania, Proceedings of the National Academy of Sciences USA 103, 2006, p. 17196-17201.
- Trinkaus 2005 E. Trinkaus, *Early modern humans*, Annual Review of Anthropology 34, 2005, p. 207-230.
- Trinkaus 2007 E. Trinkaus, European early modern humans and the fate of the Neandertals, Proceedings of the

National Academy of Sciences USA 104, 2007, p. 7367-7372.

- Trinkaus 2008 E. Trinkaus, *Behavioral implications of the Muierii 1 early modern human scapula*, Annuaire Roumain d'Anthropologie 45, p. 27–41.
- Vinogradov 1968 A. P. Vinogradov, A. L. Devirts, E. I Dobkina, N. G. Markova, *Radiocarbon dating in the Vernadsky Institute V*, Radiocarbon 10, 1968, p. 454-464.
- Wild et al. 2000 E. M. Wild, K. A. Arlamovsky, R. Golser,
 W. Kutschera, A. Priller, S. Puchegger, W. Rom, P. Steier,
 W. Vycudilik, ¹⁴C dating with the bomb peak: An application to forensic medicine. Nuclear Instruments and Methods in Physics Research 172B, 2000, p. 944-950.